

Identification of Measurement Needs to Prevent Childhood Obesity in High-Risk Populations and Environments



Kathryn E. Foti, PhD, MPH,¹ Crystal L. Perez, MPH,² Emily A. Knapp, PhD, MHS,¹ Anna Y. Kharmats, PhD,³ Amanda S. Sharfman, MS, MPH,⁴ S. Sonia Arteaga, PhD,⁵ Latetia V. Moore, PhD, MPH,⁶ Wendy L. Bennett, MD, MPH^{7,8,9}

Introduction: Children at highest obesity risk include those from certain racial/ethnic groups, from low-income families, with disabilities, or living in high-risk communities. However, a 2013 review of the National Collaborative for Childhood Obesity Research Measures Registry identified few measures focused on children at highest obesity risk. The objective is to (1) identify individual and environmental measures of diet and physical activity added to the Measures Registry since 2013 used among high-risk populations or settings and (2) describe methods for their development, adaptation, or validation.

Methods: Investigators screened references in the Measures Registry from January 2013 to September 2017 ($n=351$) and abstracted information about individual and environmental measures developed for, adapted for, or applied to high-risk populations or settings, including measure type, study population, adaptation and validation methods, and psychometric properties.

Results: A total of 38 measures met inclusion criteria. Of these, 30 assessed individual dietary ($n=25$) or physical activity ($n=13$) behaviors, and 11 assessed the food ($n=8$) or physical activity ($n=7$) environment. Of those, 17 measures were developed for, 9 were applied to (i.e., developed in a general population and used without modification), and 12 were adapted (i.e., modified) for high-risk populations. Few measures were used in certain racial/ethnic groups (i.e., American Indian/Alaska Native, Hawaiian/Pacific Islander, and Asian), children with disabilities, and rural (versus urban) communities.

Conclusions: Since 2013, a total of 38 measures were added to the Measures Registry that were used in high-risk populations. However, many of the previously identified gaps in population coverage remain. Rigorous, community-engaged methodologic research may help researchers better adapt and validate measures for high-risk populations.

Am J Prev Med 2020;59(5):746–754. © 2020 American Journal of Preventive Medicine. Published by Elsevier Inc. All rights reserved.

From the ¹Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland; ²Department of Health Policy and Management, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland; ³Department of International Health, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland; ⁴FHI 360, Washington, District of Columbia; ⁵Environmental influences on Child Health Outcomes, Office of the Director, NIH, formerly at the Division of Cardiovascular Diseases, National Heart, Lung, and Blood Institute, NIH, Bethesda, Maryland; ⁶Division of Nutrition, Physical Activity, and Obesity, Centers for Disease Control and Prevention, Atlanta, Georgia; ⁷Department of Medicine, Johns Hopkins University School of Medicine,

Baltimore, Maryland; ⁸Department of Epidemiology and Population, Johns Hopkins University School of Medicine, Baltimore, Maryland; and ⁹Department of Family and Reproductive Health, Johns Hopkins University School of Medicine, Baltimore, Maryland

Address correspondence to: Wendy L. Bennett, MD, MPH, Department of Medicine, Johns Hopkins University School of Medicine, 2024 E Monument Street, Baltimore MD 21205.

E-mail: wendy.bennett@jhmi.edu.

0749-3797/\$36.00

<https://doi.org/10.1016/j.amepre.2020.05.012>

INTRODUCTION

In the U.S., 18.5% of children aged 2–19 years have obesity.¹ Childhood obesity is a broad public health concern and leading health equity issue. Obesity prevalence is higher among Hispanic (25.8%) and non-Hispanic Black (22.0%) than White (14.1%) and Asian (11.0%) children.¹ Obesity prevalence is also elevated among children living in families with low household income or where the head of household has low educational attainment.² In addition, obesity prevalence is higher among children in rural than urban areas.³ Although there is less rigorous information, data suggest children with intellectual, developmental, or physical disabilities have a 27%–59% higher risk of obesity than those without disabilities.⁴

There is a need to develop, adapt, and validate measures for children and families at high risk for obesity to accurately assess risk factors and evaluate interventions. Most available measures are developed for general or lower-risk populations and may require modification to be valid among high-risk populations.⁵ Populations at high risk for obesity may differ from lower-risk populations in important and interrelated ways, including historical, environmental, and social contexts; literacy level or spoken language; and cultural and psychosocial perspectives on diet, physical activity, and weight control.^{6,7} In addition, they may have differential access to obesity prevention and treatment interventions.

The National Collaborative on Childhood Obesity Research (NCCOR) Measures Registry was launched in 2011 to improve the quality of research regarding dietary and physical activity behaviors and related environments, contribute to standardization across studies, and better inform policies and programs to promote the health of children.^{8,9} The Registry is a searchable database of individual and environmental dietary and physical activity measures relevant to childhood obesity research.⁹ In 2013, an Institute of Medicine (IOM) report, “Evaluating Obesity Prevention Efforts: A Plan for Measuring Progress,”¹⁰ reviewed measures in the Registry used among high-risk populations. The report focused on environmental-level measures and identified 174 (of 893) measures used among high-risk populations, but a paucity specifically developed or adapted for high-risk populations.¹⁰

The study objective is to update and expand on the previous IOM review by identifying and characterizing individual and environmental measures of diet and physical activity used among high-risk populations added to the NCCOR Measures Registry since 2013. To accomplish this goal, the authors (1) identify individual and environmental measures of diet and physical activity

used among high-risk populations or settings and (2) abstract information about their development, adaptation, or validation.

METHODS

Study Sample

The definition for high-risk populations was similar to the 2013 IOM report¹⁰ and modified based on the input from the NCCOR work group. The authors defined high-risk populations as children (aged 0–18 years) and their families at high risk for obesity or residing in communities where the risk of obesity and related comorbidities may be highest. Individual and community factors related to increased risk of obesity include race/ethnicity, disability, education/income, urbanicity, and region of the country.

NCCOR is a partnership of the 4 leading funders of childhood obesity research: the Centers for Disease Control and Prevention, NIH, Robert Wood Johnson Foundation, and U.S. Department of Agriculture. All NCCOR projects are informed by a work group of staff at the 4 agencies. The work group for this project met monthly and provided input throughout.

The NCCOR Measures Registry contains measures relevant to childhood obesity identified from literature searches of English-language articles using approximately 500 search terms. Additional details on the development of the Registry have been published previously.^{8,11} The search is updated periodically, most recently to include articles published through September 2017. The Registry currently contains nearly 1,400 measures, organized in 4 domains: individual dietary behaviors, individual physical activity behaviors, food environment, or physical activity environment. Each measure's entry contains information on how to use the measure and its validity and reliability. Examples of measures include questionnaires, logs, electronic devices, and methods for direct observation.

Measures

Investigators searched articles added to the NCCOR Measures Registry from January 2013 through September 2017 ($n=351$).⁹ They uploaded all articles into DistillerSR (Evidence Partners) for screening and data abstraction. Two trained investigators (KF and EK) independently performed title and abstract screening. Articles were included for full-text review if (1) they reported measures of 1 or more domains in high-risk populations or settings or (2) it was unclear and full-text review was needed to determine eligibility. Studies that did not include a high-risk population or setting were excluded. In addition, authors excluded studies that were (1) not conducted among children or settings applicable to children (e.g., schools or home), (2) conducted outside the U.S. (it was unclear whether high-risk populations in other countries were generalizable to the U.S.), (3) not in English, (4) published before 2013, (5) not original research, or (6) unavailable in full-text form (Figure 1).

With guidance from the NCCOR working group, investigators developed a data abstraction form (Appendix 1, available online) in the DistillerSR database. The form included data elements contained in the NCCOR Measures Registry, the 2013 IOM report,¹⁰ and related to adaptation and validation methods. The following data elements included in the NCCOR Measures Registry were abstracted: domain, measure type, study location, participant ages,




Figure 1. PRISMA diagram. NCCOR, National Collaborative on Childhood Obesity Research.

race/ethnicity, and psychometric properties of the measure. This study used similar categories as the IOM report¹⁰ to characterize sociocultural influences and SES of the study population or setting; for example, the authors abstracted information about whether the study population was described by country of origin, language proficiency, level of education, or income. They additionally abstracted whether studies included or focused on lesbian, gay, bisexual, transgender, and questioning (LGBTQ+) populations.

Investigators abstracted information about whether the measure was developed for a high-risk population, applied to a high-risk population (i.e., developed in a general population and used without modification), or adapted (i.e., modified) for use among a high-risk population; to describe what aspects of the instrument were modified; and methods for adaptation and validation. When reported by the authors, methodologic considerations for measurement among high-risk populations or settings were abstracted and summarized. The codebook is available in [Appendix 2](#) online.

For quality assurance, 2 trained reviewers (KF and CP) independently reviewed a random sample of articles and compared data abstraction. In addition, the senior author (WB) reviewed 20% of the articles to assess completeness and accuracy.

Statistical Analysis

The authors summarized the number of measures within each domain, that is, individual dietary behaviors, individual physical

activity behaviors, the food environment, and the physical activity environment. Measures that assessed multiple domains were counted in each relevant domain. The number of measures used among high-risk populations of interest were counted by domain. The authors summarized the types of measures in each domain.

The authors counted the number of measures by domain that were developed for, applied to, or adapted for high-risk populations; these categories were mutually exclusive. For measures adapted for high-risk populations, they summarized methods for adaptation and validation and how the content was modified from the original instrument.

RESULTS

A total of 38 measures from the NCCOR Measures Registry met inclusion criteria ([Appendices 3 and 4](#), available online). Of these, 30 measures assessed individual behaviors; 25 assessed individual dietary behaviors and 13 assessed individual physical activity behaviors (8 assessed both). A total of 11 measures assessed environmental determinants of obesity; 8 assessed the food environment and 7 assessed the physical activity environment (4 assessed both) ([Table 1](#)).

Table 1. Summary of the Number of NCCOR Measures Registry Tools Identified Since 2013 Targeting Obesity Prevention Efforts for High-Risk Populations and Settings

High-risk population or setting	Individual behavior measures (n=30 measures)			Environmental measures (n=11 measures)		
	Dietary behavior (n=25) n (%)	Physical activity behavior (n=13) n (%)	By subpopulation n (%)	Food environment (n=8) n (%)	Physical activity environment (n=7) n (%)	By setting n (%)
Racial/ethnic group						
African American	16 (64)	6 (46)	18 (60)	3 (38)	3 (43)	5 (45)
American Indian/ Alaska Native	5 (20)	1 (8)	5 (17)	2 (25)	2 (29)	2 (18)
Hispanic	19 (76)	9 (69)	23 (77)	4 (50)	4 (57)	6 (55)
Hawaiian/Pacific Islander	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Asian	6 (24)	2 (15)	6 (20)	1 (13)	1 (14)	1 (9)
White	11 (44)	4 (31)	12 (40)	3 (38)	3 (43)	5 (45)
Other	10 (40)	5 (38)	12 (40)	4 (50)	2 (29)	5 (45)
Not reported	1 (4)	0 (0)	1 (3)	2 (25)	2 (29)	3 (27)
Disability/special healthcare needs	1 (4)	0 (0)	1 (3)	0 (0)	0 (0)	0 (0)
Geographic location						
Metro/urban	16 (64)	9 (69)	21 (70)	5 (63)	5 (71)	8 (73)
Small town/rural	4 (16)	2 (15)	4 (13)	4 (50)	4 (57)	5 (45)
Social influences						
Low income/SES	15 (60)	6 (46)	17 (57)	6 (75)	5 (71)	8 (73)
Low education	6 (24)	4 (31)	8 (27)	2 (25)	1 (14)	2 (18)
Language proficiency	4 (16)	3 (23)	6 (20)	1 (13)	0 (0)	1 (9)
Acculturation	3 (12)	2 (15)	4 (13)	0 (0)	0 (0)	0 (0)
Foreign born	3 (12)	2 (15)	4 (13)	1 (13)	0 (0)	0 (0)
Living and working conditions	3 (12)	2 (15)	4 (13)	2 (25)	2 (29)	2 (18)
Racial/ethnic composition of community	6 (24)	2 (15)	6 (20)	2 (25)	4 (57)	4 (36)
Measure type						
24-hour dietary recall	2 (8)	N/A	2 (7)	N/A	N/A	N/A
Food frequency questionnaire	4 (16)	N/A	4 (13)	N/A	N/A	N/A
Other questionnaire	16 (64)	11 (85)	19 (63)	5 (63)	3 (43)	5 (45)
Record or log	0 (0)	1 (8)	1 (3)	0 (0)	0 (0)	0 (0)
Electronic monitor	0 (0)	1 (8)	1 (3)	0 (0)	0 (0)	0 (0)
Interview	1 (4)	0 (0)	1 (3)	0 (0)	0 (0)	0 (0)
Behavioral observation	0 (0)	0 (0)	0 (0)	N/A	N/A	N/A
Environmental observation	N/A	N/A	N/A	2 (25)	3 (43)	5 (45)
GIS	N/A	N/A	N/A	1 (13)	1 (14)	1 (9)
Other	2 (12)	0 (0)	2 (7)	0 (0)	0 (0)	0 (0)
Use in high-risk population						
Applied to high-risk population	8 (32)	3 (23)	9 (30)	1 (13)	1 (14)	2 (18)
Developed for high-risk population	10 (40)	6 (46)	13 (43)	4 (50)	3 (43)	5 (45)
Adapted for high-risk population	7 (28)	4 (31)	8 (27)	3 (38)	3 (43)	4 (36)

NCCOR, National Collaborative on Childhood Obesity Research.

November 2020

Table 2. Considerations for Developing or Adapting Measures for High-Risk Populations as Described in Articles Included in the National Collaborative on Childhood Obesity Research Measures Registry

Measure type	Considerations
Individual behavior measures	
Food frequency or other questionnaire	<p>Include culturally relevant foods.^{12,23–25}</p> <p>Availability of foods will vary by region. Consider local food availability and where food is sourced. For example, in one study conducted in Puerto Rico, most foods consumed were imported from the Continental U.S.²⁶</p> <p>Level of acculturation and education may influence the difficulty in responding to a food frequency questionnaire.²³</p> <p>Describe foods and beverages in ways that are familiar to certain cultural groups to help improve validity.¹²</p> <p>Consider cultural differences in perception of healthfulness of sugar-sweetened beverages (e.g., sport drinks) and culturally relevant sweetened drinks (e.g., aguas frescas, which contain sugar, fruit, and water). Misconceptions have been reported among Hispanic youth.¹²</p> <p>Systematic biases (e.g., by personal characteristics such as body weight, social or cultural desirability, acculturation level, or literacy level) may influence the reporting of dietary intake, with a larger variance and reduced correlations with true intake.^{12,27}</p>
High-risk population	
Acculturation status	First-, second-, and third-generation immigrants may have different health beliefs and behaviors. The influence of acculturation may also vary by country of origin. ²³
Language proficiency	Respondents who choose to complete a measure in English may systematically differ from those who choose to complete it in another language. ¹³
Food insecure	Capture the child's perspective as children may experience food insecurity differently from their parents or caregivers. Measures of child and adult food insecurity may be more appropriate than a single adult or household measure. ²⁸
Environmental measures	
Environmental observation (home)	<p>Include culturally relevant foods, particularly for racial/ethnic minorities and recent immigrants.^{14,24}</p> <p>Timing of grocery shopping will affect the foods available in the home.¹⁴</p> <p>Include activities available for families with socioeconomic, racial, and ethnic diversity.²⁴</p>
Population or setting	
Rural	<p>Many rural residents have Post Office Box mailing addresses. This will affect the validity of GIS measures of the food and physical activity environment based on participant address in rural settings.²⁹</p> <p>Season and rurality may impact food availability. Certain fruits and vegetables may not be available in very rural areas nor in specific climates.³⁰</p>
Community safety	Safety may influence physical activity. In one study, American Indian children living on a reservation reported feeling unsafe when using their community bike path and a lack of resources to engage in physical activity. It is important to consider such barriers when developing measures and interventions. ³¹

Individual behavior measures were most commonly used among Hispanic ($n=23$, 77%) and African American ($n=18$, 60%) populations. Six studies included Asian populations (20%), 5 (17%) included American Indian/Alaska Native populations, and none included Hawaiian/Pacific Islander populations. One measure was used among children with autism or other special healthcare needs. There were no measures used among LGBTQ+ populations. Measures were more commonly used among populations living in metropolitan or urban areas ($n=21$, 70%) than small towns or rural areas ($n=4$, 13%). A total of 17 measures assessed individual behaviors among low-income or low-SES populations (57%). Measures more commonly assessed behaviors among

children aged 6–11 years ($n=16$, 53%) and 2–5 years ($n=15$, 50%) than those aged 12–18 years ($n=7$, 23%) or <2 years ($n=2$, 7%). Questionnaires were the most frequently identified measure type ($n=19$, 63%).

There were 13 individual-level measures (43%) developed for high-risk populations, 9 (30%) applied to high-risk populations (i.e., developed in a general population and used without modification), and 8 (27%) adapted (i.e., modified) for use among high-risk populations. Authors described several considerations for measuring individual dietary and physical activity behaviors among high-risk populations (Table 2). Cultural and linguistic adaptations were the most common forms of adaptation. Most often, researchers modified dietary measures to be

more culturally appropriate ($n=6$). Focus group discussions, other qualitative, and mixed-methods approaches were often used for these purposes.

For example, one study adapted a validated adult Beverage Intake Questionnaire (BEVQ-15) for use among Hispanic preschool children aged 3–5 years.¹² Researchers conducted 20 semistructured interviews with Hispanic mothers to identify relevant beverages from the original instrument, add suggested beverages, and adapt serving sizes for young children. The adapted instrument was piloted ($n=5$ mothers) and refined based on feedback on the questions, format, and mode of administration. In a validation/reliability study, 109 mothers completed the BEVQ for preschoolers, or BEVQ-PS, which was compared with a 4-day food intake record. Test–retest reliability was assessed over a 6 to 9-day period. The authors found that sugar-sweetened beverages, whole milk, and water met validity and reliability criteria, but modifications may be needed to accurately assess total beverage intake.¹²

In another study, researchers developed the Preschooler Physical Activity Parenting Practices instrument¹³ for use among Latino parents. The instrument was developed based on formative qualitative research using the Nominal Group Technique. Latino parents were asked what they do to encourage or discourage physical activity. Responses were ordered and grouped into parenting factors based on the literature. The instrument was translated into Spanish and back-translated into English. Conceptual and cultural, rather than linguistic, equivalence was prioritized when there were differences between the original and back-translated survey. Researchers also conducted cognitive interviews with 5 English-speaking and 5 Spanish-speaking participants to refine survey items. Although the instrument showed moderate to excellent test–retest reliability and acceptable internal consistency, only certain subscales were significantly correlated with accelerometer-measured child physical activity. The authors noted that, for some subscales regarding parenting practices that discouraged physical activity, the Cronbach's α was lower for the Spanish- than English-language version of the instrument. Spanish-speaking participants had lower levels of education, which may have confounded this observation.

Of the 11 environmental-level measures, there were 6 (55%) used among Hispanic populations, 5 (45%) in African American populations, 2 in American Indian/Alaska Native populations, 1 in an Asian population, and none in Hawaiian/Pacific Islander populations. No environmental-level measures were used among children with disabilities or special healthcare needs. Measures were more commonly used in metropolitan or urban settings ($n=8$, 73%) than small towns or rural areas ($n=5$, 45%). Eight measures were used in low-income or low-

SES populations or settings (73%). Environmental observation ($n=5$, 45%) and questionnaires ($n=5$, 45%) were the most common measure types; there was one GIS measure.

Five environmental measures (45%) were developed for high-risk settings, 2 (18%) were applied to high-risk settings (i.e., developed in general settings and used without modification), and 4 (36%) were adapted for high-risk settings (i.e., modified). One measure that was adapted was a home food inventory for low-income Spanish- and Somali-speaking families with preschool-aged children.¹⁴ Focus groups were conducted with 5 Spanish-speaking and 5 Somali-speaking individuals with English-language skills to update an existing home food inventory. The updated inventory was translated into Spanish and Somali. The inventory was validated comparing responses of a trained staff member with those from 15 Spanish-speaking and 15 Somali-speaking parents. All validity indices were in an acceptable range, except for specific items such as whole wheat bread, possibly because of language or literacy barriers combined with poor understanding of nutrition labels among the general population. The authors concluded that the tool is a valid measure among Spanish and Somali households and should be validated in other populations.

The Texas Childhood Obesity Prevention Policy Evaluation School Environmental Audit Tool was developed to assess the safety and walkability of school environments.¹⁵ The tool, developed from a conceptual framework, includes street, school site, and map audits. It was pre-tested in 1 urban, suburban, and rural elementary school; refined; and tested again. Two trained auditors then visited 12 elementary schools (4 urban, 4 suburban, and 4 rural, including 2 high- and 2 low-income schools in each area) to assess inter-rater, test–retest, and peak versus off-peak hour reliability. Test–retest and peak versus off-peak reliability were highest among rural schools. Inter-rater reliability was highest at urban schools and lowest at rural schools for perceptual qualities (e.g., safety and attractiveness), likely because of heterogeneity in rural environments; inter-rater reliability for objective items was excellent for all settings. The authors concluded that, with proper training to reduce inter-rater differences, this tool can assess school environments reliably across settings for surveillance, research, and policy evaluation.

DISCUSSION

Since the 2013 IOM review of the NCCOR Measures Registry, an increasing number of measures are being developed or adapted for high-risk populations. However, measurement gaps for specific populations and settings were similar to those identified in the previous

report. Although a large proportion of measures identified in this review were used among African American and Hispanic individuals and communities, fewer were used among Asians, American Indians/Alaska Natives, or Hawaiian/Pacific Islanders, despite the high prevalence of overweight and obesity among American Indian and Hawaiian/Pacific Islander children¹⁶ and substantial, understudied heterogeneity by Asian ethnicity.^{17,18} Although country of origin and acculturation may influence knowledge, beliefs, and behaviors related to diet and physical activity, these factors are rarely assessed; additional tailoring of measures may be needed. As in the 2013 report, this study identified a need for measures for children with disabilities and special healthcare needs as only 1 relevant measure was found. In addition, there is a critical gap in measures used among LGBTQ+ populations, as none were identified in the 2013 report or this study. More measures were applied in urban than rural settings, similar to the previous report. Few studies reported sufficient detail to precisely classify geographic areas. The choice of rural/urban classification scheme may affect the validity of certain measures.¹⁹

This study found that most measures used in high-risk populations were questionnaires. Although there are advantages with respect to participant burden and ease of administration, questionnaires have limitations such as the potential for recall bias or reporting errors. In addition, language or literacy barriers among certain subgroups could further affect validity. The type of instrument and mode of administration may be particularly important considerations when engaging high-risk populations.

There are several challenges related to measurement in high-risk populations to be addressed. First is to promote the use of best practices for adaptation and standard validation procedures.²⁰ Few studies reported details about adaptation methods used, and the quality of such studies varied. Further, a number of studies did not discuss differences in the validity of subscales within a measure or differences across populations (if applicable) nor the limitations on the context in which the measure could provide useful information.²¹ The authors see a need for rigorous methodologic research and to increase dissemination of adaptation and validation studies, which may not necessarily be published in the literature. A second challenge is to balance the tension between tailoring measures for specific groups and using standardized measures to facilitate comparison across populations.²¹ Researchers will need to consider the trade-offs and select measures appropriate for the purpose of their work. Third is intersectionality. Disparities in childhood obesity are rarely explained by a single factor.²² Characteristics used to define high-risk populations often co-occur and interact. Researchers will need

to consider how their intersection may influence measurement and the implications for practice and policy responses. A final challenge is to increase community engagement. Few studies described how community members were engaged in measure development or adaptation, which ultimately may affect validity. Community members' perspectives are critical to measure what matters and understand how to measure it.²⁰ Researchers conducting community-engaged studies can advance the field by documenting and sharing best practices and lessons learned.

This study has several limitations. First, it focused on measures included in the NCCOR Measures Registry and may have missed other studies using these measures that were not captured. However, the NCCOR search strategy is rigorous.¹¹ In addition, individual behavior measures included in the NCCOR Measures Registry are required to be previously validated; there may be other tools used in practice that are not included in the Registry. Second, it relied on study descriptions about setting and urbanicity/rurality. Third, it relied on the authors' descriptions of adaptation and validation methods and considerations for measures used among specific high-risk groups, which were often only briefly reported.

Recognizing limited progress, NCCOR has taken steps to identify measurement priorities to address gaps related to children in high-risk populations or settings. In September 2019, NCCOR held a 2-day workshop, titled "Advancing Measurement for High-Risk Populations and Communities Related to Childhood Obesity," with a goal of illustrating current challenges, discussing best practices to adapt and develop measures, and developing recommendations to address gaps in the field. The workshop convened more than 20 experts in the measurement of high-risk populations. Recommendations from the workshop will be shared on the NCCOR website, www.nccor.org/measurement-workshop-series/.

CONCLUSIONS

To reduce disparities in childhood obesity, it is necessary to measure individual behaviors and environmental factors in the socioeconomic and sociocultural contexts in which they occur. This report provides an overview of the current state of measures available for use in high-risk populations. Although there has been an increase in measures used among high-risk populations since 2013, there are certain populations and settings for which major gaps remain. It is also important to understand whether methodological choices related to the development and adaptation of measures for high-risk populations achieve the goal of accurately measuring constructs of interest. These issues may be especially salient among

high-risk populations and in disadvantaged neighborhoods, as well as in rural areas. Addressing gaps in the availability of validated tools and measures and improving the quality of measurement can help practitioners understand and address risk factors for obesity among high-risk children and their families.

ACKNOWLEDGMENTS

This work was funded by The JPB Foundation (Grant Number 949).

Dr. Arteaga is now a Program Official of the Environmental influences on Child Health Outcomes Program Office, Office of the Director, National Institutes of Health, Bethesda, MD.

No financial disclosures were reported by the authors of this paper.

The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention, NIH, or HHS.

SUPPLEMENTAL MATERIAL

Supplemental materials associated with this article can be found in the online version at <https://doi.org/10.1016/j.amepre.2020.05.012>.

REFERENCES

- Hales CM, Carroll MD, Fryar CD, Ogden CL. Prevalence of obesity among adults and youth: United States, 2015–2016. *NCHS Data Brief*. 2017;(288):1–8. www.cdc.gov/nchs/data/databriefs/db288.pdf. Accessed July 17, 2020.
- Ogden CL, Carroll MD, Fakhouri TH, et al. Prevalence of obesity among youths by household income and education level of head of household - United States 2011–2014. *MMWR Morb Mortal Wkly Rep*. 2018;67(6):186–189. <https://doi.org/10.15585/mmwr.mm6706a3>.
- Johnson JA 3rd, Johnson AM. Urban–rural differences in childhood and adolescent obesity in the United States: a systematic review and meta-analysis. *Child Obes*. 2015;11(3):233–241. <https://doi.org/10.1089/chi.2014.0085>.
- Bandini L, Danielson M, Esposito LE, et al. Obesity in children with developmental and/or physical disabilities. *Disabil Health J*. 2015;8(3):309–316. <https://doi.org/10.1016/j.dhjo.2015.04.005>.
- Stewart AL, Thrasher AD, Goldberg J, Shea JA. A framework for understanding modifications to measures for diverse populations. *J Aging Health*. 2012;24(6):992–1017. <https://doi.org/10.1177/0898264312440321>.
- Kumanyika S. *Getting to equity in obesity prevention: a new framework*. Washington, DC: National Academy of Medicine, 2017. <https://doi.org/10.31478/201701c>. Published January 18, 2017. Accessed August 21, 2019.
- Kumanyika S. The sociocultural context for obesity prevention and treatment in children and adolescents: Influences of ethnicity and gender. In: Freemark M, ed. *Pediatric Obesity. Contemporary Endocrinology*. Humana Press, Cham; 2018:695–713. https://doi.org/10.1007/978-3-319-68192-4_40.
- McKinnon RA, Reedy J, Berrigan D, Krebs-Smith SM, NCCOR Catalogue and Registry Working Groups. The National Collaborative on Childhood Obesity Research Catalogue of Surveillance Systems and Measures Registry: new tools to spur innovation and increase productivity in childhood obesity research. *Am J Prev Med*. 2012;42(4):433–435. <https://doi.org/10.1016/j.amepre.2012.01.004>.
- Measures Registry. National Collaborative on Childhood Obesity Research. <https://tools.nccor.org/measures>. Accessed October 21, 2019.
- Institute of Medicine. *Evaluating obesity prevention efforts: a plan for measuring progress*. Washington, DC: National Academies Press, 2013. <https://doi.org/10.17226/18334>. Published August 2, 2013. Accessed June 29, 2018.
- Finkelstein D, Fox MK. *Development of the database for the National Collaborative on Childhood Obesity Research Measures Registry: final report*. Princeton, NJ: Mathematica Policy Research, Inc.; January 11, 2011. www.nccor.org/downloads/NCCOR%20MPR%20Report%20Final.pdf.
- Lora KR, Davy B, Hedrick V, Ferris AM, Anderson MP, Wakefield D. Assessing initial validity and reliability of a beverage intake questionnaire in Hispanic preschool-aged children. *J Acad Nutr Diet*. 2016;116(12):1951–1960. <https://doi.org/10.1016/j.jand.2016.06.376>.
- O'Connor TM, Cerin E, Hughes SO, et al. Psychometrics of the Preschooler Physical Activity Parenting Practices instrument among a Latino sample. *Int J Behav Nutr Phys Act*. 2014;11:3. <https://doi.org/10.1186/1479-5868-11-3>.
- Hearst MO, Fulkerson JA, Parke M, Martin L. Validation of a home food inventory among low-income Spanish- and Somali-speaking families. *Public Health Nutr*. 2013;16(7):1151–1158. <https://doi.org/10.1017/S1368980012004508>.
- Lee C, Kim HJ, Dowdy DM, Hoelscher DM, Ory MG. TCOPPE school environmental audit tool: assessing safety and walkability of school environments. *J Phys Act Health*. 2013;10(7):949–960. <https://doi.org/10.1123/jpah.10.7.949>.
- Pan L, Freedman DS, Park S, Galuska DA, Potter A, Blanck HM. Changes in obesity among U.S. children aged 2 through 4 years enrolled in WIC during 2010–2016. *JAMA*. 2019;321(23):2364–2366. <https://doi.org/10.1001/jama.2019.5051>.
- Asian & Pacific Islander American Health Forum. Obesity and overweight among Asian American children and adolescents. www.apiahf.org/wp-content/uploads/2016/04/2016.04.28_OBESITY-AND-OVERWEIGHT-AMONG-AA-CHILDREN-AND-ADOLESCENTS_Data-Brief-1.pdf. Published 2016. Accessed January 16, 2020.
- Quader ZS, Gazmararian JA, McCullough LE. Obesity and understudied minority children: existing challenges and opportunities in epidemiology. *BMC Pediatr*. 2019;19(1):103. <https://doi.org/10.1186/s12887-019-1484-9>.
- National Academies of Sciences, Engineering, and Medicine. *Rationalizing rural area classifications for the Economic Research Service: a workshop summary*. Washington, DC: National Academies Press, 2016. <https://doi.org/10.17226/21843>. Published 2016. Accessed November 25, 2019.
- Bender MS, Clark MJ. Cultural adaptation for ethnic diversity: a review of obesity interventions for preschool children. *Calif J Health Promot*. 2011;9(2):40. <https://doi.org/10.32398/cjhp.v9i2.1435>.
- Frongillo EA, Baranowski T, Subar AF, Toozee JA, Kirkpatrick SI. Establishing validity and cross-context equivalence of measures and indicators. *J Acad Nutr Diet*. 2019;119(11):1817–1830. <https://doi.org/10.1016/j.jand.2018.09.005>.
- López N, Gadsen VL. *Health inequities, social determinants, and intersectionality*. Washington, DC: National Academy of Medicine, 2016. <https://doi.org/10.31478/201612a>. Published December 5, 2016. Accessed November 25, 2019.
- Park SH, Park CG, McCreary L, Norr KF. Cognitive interviews for validating the Family Nutrition Physical Activity instrument for Korean-American families with young children. *J Pediatr Nurs*. 2017;36:1–6. <https://doi.org/10.1016/j.pedn.2017.04.014>.

24. Boles RE, Burdell A, Johnson SL, Gavin WJ, Davies PL, Bellows LL. Home food and activity assessment. Development and validation of an instrument for diverse families of young children. *Appetite*. 2014; 80:23–27. <https://doi.org/10.1016/j.appet.2014.04.026>.
25. Hartman TJ, Elliott AJ, Angal J, et al. Relative validation of a short questionnaire to assess the dietary habits of pregnant American Indian women. *Food Sci Nutr*. 2017;5(3):625–632. <https://doi.org/10.1002/fsn3.440>.
26. Palacios C, Rivas-Tumanyan S, Santiago-Rodríguez EJ, et al. A semi-quantitative food frequency questionnaire validated in Hispanic infants and toddlers aged 0 to 24 months. *J Acad Nutr Diet*. 2017;117(4):526–535. e9. <https://doi.org/10.1016/j.jand.2016.12.010>.
27. Matthews-Ewald MR, Posada A, Wiesner M, Olvera N. An exploratory factor analysis of the Parenting Strategies for Eating and Physical Activity Scale (PEAS) for use in Hispanic mothers of adolescent and preadolescent daughters with overweight. *Eat Behav*. 2015;19:193–199. <https://doi.org/10.1016/j.eatbeh.2015.10.001>.
28. Baxter SD, Smith AF, Hitchcock DB, et al. Test–retest reliability of the National Health and Nutrition Examination Survey’s 5-item Food Insecurity Questionnaire completed by fourth-grade children. *J Nutr Educ Behav*. 2015;47(5):459–464. e1. <https://doi.org/10.1016/j.jneb.2015.06.006>.
29. Lee JE, Sung JH, Malouhi M. Statistical validation of a web-based GIS application and its applicability to cardiovascular-related studies. *Int J Environ Res Public Health*. 2015;13(1):ijerph13010002. <https://doi.org/10.3390/ijerph13010002>.
30. Byker Shanks C, Jilcott Pitts S, Gustafson A. Development and validation of a farmers’ market audit tool in rural and urban communities. *Health Promot Pract*. 2015;16(6):859–866. <https://doi.org/10.1177/1524839915597899>.
31. Arcan C, Hannan PJ, Fulkerson JA, et al. Associations of home food availability, dietary intake, screen time and physical activity with BMI in young American-Indian children. *Public Health Nutr*. 2013;16(1):146–155. <https://doi.org/10.1017/S136898001200033X>.