

Surgical Site Infection Prevention

A Review

Jessica L. Seidelman, MD, MPH; Christopher R. Mantyh, MD; Deverick J. Anderson, MD, MPH



IMPORTANCE Approximately 0.5% to 3% of patients undergoing surgery will experience infection at or adjacent to the surgical incision site. Compared with patients undergoing surgery who do not have a surgical site infection, those with a surgical site infection are hospitalized approximately 7 to 11 days longer.

OBSERVATIONS Most surgical site infections can be prevented if appropriate strategies are implemented. These infections are typically caused when bacteria from the patient's endogenous flora are inoculated into the surgical site at the time of surgery. Development of an infection depends on various factors such as the health of the patient's immune system, presence of foreign material, degree of bacterial wound contamination, and use of antibiotic prophylaxis. Although numerous strategies are recommended by international organizations to decrease surgical site infection, only 6 general strategies are supported by randomized trials. Interventions that are associated with lower rates of infection include avoiding razors for hair removal (4.4% with razors vs 2.5% with clippers); decolonization with intranasal antistaphylococcal agents and antistaphylococcal skin antiseptics for high-risk procedures (0.8% with decolonization vs 2% without); use of chlorhexidine gluconate and alcohol-based skin preparation (4.0% with chlorhexidine gluconate plus alcohol vs 6.5% with povidone iodine plus alcohol); maintaining normothermia with active warming such as warmed intravenous fluids, skin warming, and warm forced air to keep the body temperature warmer than 36 °C (4.7% with active warming vs 13% without); perioperative glycemic control (9.4% with glucose <150 mg/dL vs 16% with glucose >150 mg/dL); and use of negative pressure wound therapy (9.7% with vs 15% without). Guidelines recommend appropriate dosing, timing, and choice of preoperative parenteral antimicrobial prophylaxis.

CONCLUSIONS AND RELEVANCE Surgical site infections affect approximately 0.5% to 3% of patients undergoing surgery and are associated with longer hospital stays than patients with no surgical site infections. Avoiding razors for hair removal, maintaining normothermia, use of chlorhexidine gluconate plus alcohol-based skin preparation agents, decolonization with intranasal antistaphylococcal agents and antistaphylococcal skin antiseptics for high-risk procedures, controlling for perioperative glucose concentrations, and using negative pressure wound therapy can reduce the rate of surgical site infections.

JAMA. 2023;329(3):244-252. doi:10.1001/jama.2022.24075

Author Affiliations: Duke Center for Antimicrobial Stewardship and Infection Prevention, Duke University School of Medicine, Durham, North Carolina (Seidelman, Anderson); Department of Surgery, Duke University School of Medicine, Durham, North Carolina (Mantyh).

Corresponding Author: Deverick J. Anderson, MD, MPH, DUMC, Box 102359, Durham, NC 27710 (deverick.anderson@duke.edu).

Section Editor: Mary McGrae McDermott, MD, Deputy Editor.

A surgical site infection is defined as infection following an operation at an incision site or adjacent to the surgical incision.¹ Infections occur in approximately 0.5% to 3% of patients undergoing surgery²⁻⁴ and are among the most prevalent health care-acquired infections.⁵⁻⁷ Surgical site infections are responsible for approximately \$3.5 billion to \$10 billion in US health care costs annually.^{8,9} Compared with patients without surgical site infections, those with them remain in the hospital approximately 7 to 11 days longer^{7,10}; 1 study involving 177 706 postsurgical patients reported that 78% were readmitted as a result of the infection.¹¹ This review summarizes current evidence-based interventions for prevention of surgical site infection that are applicable to the majority of operations (Box).

Methods

We searched PubMed, Google Scholar, and the Cochrane database for English-language studies of pathogenesis, clinical presentation, and prevention of surgical site infections published from January 1, 2016, when guidelines were most recently published by the World Health Organization, to September 15, 2022. In addition, we manually searched the references of selected articles for additional relevant publications. We prioritized randomized trials, systematic reviews, meta-analyses, clinical practice guidelines, and articles pertinent to general medical readership. Of 94 studies identified, 69 were included, consisting of 14 randomized trials, 19

systematic reviews, 12 meta-analyses, 4 clinical practice guidelines, 17 cohort studies, and 3 cross-sectional studies.

Discussion and Observations

Pathophysiology

Surgical site infection acquisition depends on several factors, namely, exposure to bacteria and the host's ability to control the inevitable bacterial contamination of the incision. They are typically caused by bacteria inoculated into the surgical site at the time of surgery. Approximately 70% to 95% are caused by the patient's endogenous flora.¹² The most common organisms are *Staphylococcus aureus*, coagulase-negative *Staphylococcus*, and *Escherichia coli*.¹³ In some patients, introduction of only 100 colony-forming units of bacteria into the surgical site can cause infection.¹⁴ However, exogenous sources of contamination during surgery such as bacteria transmitted from surgical personnel or heater-cooler units can also lead to infections.

Pathogens that cause infection vary by surgical location. The most common pathogens are components of skin flora such as *S aureus* and *Streptococcus* species. In contrast, infections following gastrointestinal procedures are typically associated with enteric organisms such as *Enterococcus* species and *E coli*.¹⁵ Overall, *S aureus* is the most common cause of infection; for example, *S aureus* was associated with 24% of nonsuperficial surgical site infections in a cohort study including 32 community hospitals in the southeastern US.⁴ Although methicillin-resistant *S aureus* (MRSA) was previously more likely to cause surgical site infections than methicillin-sensitive *S aureus* (MSSA), the rate of MSSA-derived infections from 2013 to 2018 was higher (0.07 per 100 procedures) than the rate of MRSA infections during the same period (0.05 per 100 procedures).⁴ MRSA surgical site infections lead to worse clinical outcomes than those caused by less resistant pathogens.¹⁰ Specifically, compared with MSSA surgical site infections, those due to MRSA were independently associated with 5.5 additional hospital days (95% CI, 1.97-9.11).¹⁰ *E coli* and *Enterococcus* species respectively cause approximately 9.5% and 5.1% of all surgical site infections.¹³

Factors Associated With Surgical Site Infection

Factors associated with surgical site infection include older age, presence of immunosuppression, obesity, diabetes, effectiveness of antimicrobial prophylaxis, surgical site tissue condition (such as the presence of foreign material), and degree of wound contamination (Table 1 and Table 2). For example, a national study of more than 387 000 patients found that for most surgery types, rates of surgical site infection were increased in patients with obesity.²¹ The rates of surgical site infection following mastectomy among 16 473 patients increased with body mass index (BMI), calculated as weight in kilograms divided by height in meters squared. Those with a BMI of 20 to 25 had a surgical site infection rate of 4.66%; BMI of more than 30 to 40, 7.06%; and BMI of more than 40, 10.58%. Similarly, after 29 603 laparoscopic cholecystectomy procedures (urgency not specified), the infection rate increased with BMI: 8.57% with a BMI of 20 to 25; 10.62% with a BMI of 30 to 40; and 17.11% with a BMI of more than 40.

Box. Commonly Asked Questions

How can the generalist clinician help in preventing surgical site infections?

The generalist can help patients improve modifiable characteristics associated with increased risk of surgical site infections, such as helping obese patients lose weight, assisting patients who have diabetes with optimal glucose control, and assisting with smoking cessation.

Is there a threshold hemoglobin A_{1c} value above which surgical site infections are more common and surgery should be delayed?

Perioperative hyperglycemia in patients with or without diabetes is associated with surgical site infections, and randomized clinical trials support perioperative glucose control as an evidence-based practice to decrease risk of surgical site infection. In contrast, there are no randomized clinical trials that have found a clear association between a specific hemoglobin A_{1c} cutoff value and surgical site infections. However, patients with higher hemoglobin A_{1c} levels will likely have higher perioperative glucose values and glucose levels that are harder to control.

What therapies can prevent a surgical site infection?

Numerous strategies are currently recommended as outlined in this review. Six are supported by randomized clinical trials: (1) do not remove hair at the surgical site unless necessary; (2) decolonization with intranasal antistaphylococcal agent and antistaphylococcal skin antiseptic prior to high-risk procedures (eg, cardiothoracic, orthopedic); (3) use a chlorhexidine gluconate-alcohol antiseptic agent for skin preparation; (4) maintain normothermia intraoperatively; (5) control perioperative glucose values between 110 and 150 mg/dL; and (6) use incisional negative pressure wound dressings.

Some of these risk factors associated with surgical site infection are modifiable, such as hyperglycemia, obesity, and tobacco use. Other factors are nonmodifiable, such as age, which must be considered when deciding on the surgical intervention for the patient.^{26,49}

Clinical Presentation

The median time to diagnosis of surgical site infection varies by procedure.⁵⁰ For example, *S aureus* infection is typically diagnosed a median of 14 days after plastic surgery, 24 days after general orthopedic surgery, and 28 days after orthopedic surgery where a prosthetic device was inserted. A surgical site infection is suspected when purulent drainage is present at the incision site or when there is evidence of an abscess involving the surgical bed. Physical examination findings such as systemic signs of infection (eg, fevers, rigors), local erythema, wound dehiscence, pain, nonpurulent drainage, or induration are the most common. However, the presence or absence of these symptoms varies depending on factors such as surgical site, host, and time from surgery to presentation. For example, fevers can be present in 14% of patients with a chronic prosthetic joint infection but up to 75.5% of patients if the etiology of the prosthetic joint infection is hematogenous.⁵¹ Articular effusion and swelling may be present in 29% to 75% of prosthetic joint infections of the knee,⁵² and delayed wound healing, wound dehiscence, or wound drainage

Table 1. Modifiable and Nonmodifiable Patient-Related Factors Associated With Surgical Site Infections

Factor	Pathophysiology
Patient-related, modifiable	
Diabetes	Hyperglycemia impairs the innate immune system and promotes glycosylation of proteins, which compromises wound healing. ¹⁶ Diabetes can lead to higher perioperative glucose levels and hyperglycemia that is more difficult to treat. ¹⁷
Immunosuppressive medications and conditions	Immunosuppressive clinical conditions or medications diminish the inflammatory phase of wound healing. ^{18,19}
Malnutrition	Malnutrition can decrease collagen synthesis, granulation formation in surgical wounds, and result in poor tissue healing. Hypoalbuminemia weakens innate immunity by prompting macrophage apoptosis and diminishing macrophage activation. Low albumin also accelerates the seepage of interstitial fluid into the surgical wound and promotes general tissue edema. ²⁰
Obesity	Adipose tissue has less blood flow, which inhibits the delivery of oxygen and antibiotics. ²¹⁻²³
Preoperative infections	Prior to elective surgery, recognize and treat all infections (even if they are distant from the surgical site). ²⁴
Tobacco use	Tobacco use causes vasoconstriction, which can progress to alterations in collagen metabolism, decreased inflammatory response, and relative ischemia. ²⁵
Patient-related, nonmodifiable	
Age	The skin's basement membrane and dermis thin with increasing age, and the skin loses its reserve of cutaneous blood vessels and nerves that diminish wound healing. ^{26,27}
History of prior skin and soft tissue infections	A history of skin and soft tissue infections may be indicative of issues with inherent immunity and propensity for infection. ²⁸
History of radiation therapy	Treatment with radiation induces underlying tissue injury and inhibits wound healing.

Table 2. Modifiable Operation-Related Factors Associated With Surgical Site Infections

Factor	Pathophysiology
Airborne contamination	Raising the amount of microorganisms in the operating room environment provides additional opportunity for surgical site infection. Most of the airborne pathogens are generated by persons in the operating room and their movements. ^{29,30}
Anticoagulation	Anticoagulants may generate continual oozing of the incision and slow wound healing. ³¹
Blood transfusions	Blood transfusions impair macrophage activity and influence infection risk. ³²
Decreased tissue oxygenation	Diminished tissue oxygenation lends itself to decreased oxidative killing by neutrophils and impaired tissue healing from depleted epithelialization, neovascularization, and collagen formation. Low oxygen settings can curtail the efficacy of perioperative antibiotics. ^{33,34}
Foreign material	Foreign material stimulates inflammation at the surgical site and raises the risk of surgical site infection. ^{35,36}
Operation length	Longer operative time is associated with higher damage to wound cells, wound contamination, and exposure to the outside environment. ³⁷
Perioperative hypothermia	Perioperative hypothermia weakens immune system protection against surgical wound contamination: vasoconstriction leads to impaired tissue perfusion and less access for key immune cells, less motility of key immune cells, and decreased scar formation. ³⁸
Postoperative hyperglycemia	Cellular functions of bactericidal activity, leukocyte adherence chemotaxis, and phagocytosis are enhanced by insulin and glycemic control, suggesting a direct relation between elevated blood glucose and cellular function deficits. ³⁹ This relationship is observed in patients with and without a diagnosis of diabetes.
Surgical technique	Wound healing is decreased by leaving behind devitalized tissues, inadvertent entry into hollow viscera, inadequate blood supply maintenance, rough manipulation of tissue, misplaced drains and sutures, and unsuitable postoperative wound care. ⁴⁰
Wound care	Wounds that remain uncovered following surgery can be contaminated, or uncontrolled drainage can diminish the integrity of the surrounding skin. ^{41,42}
Wound contamination from patient's own flora	Wound classification delineates the degree of contamination of a surgical wound at the time of the operation. ⁴³ Skin preparation and perioperative antibiotic administration reduce but do not eliminate the introduction of microorganisms at the surgical site. ^{44,45} Shaving leads to microscopic cuts in the skin that can become niduses for bacteria to multiply. ⁴⁰ Without appropriate drapes and barrier devices, bacteria from hair follicles and deeper skin layers can recolonize the surgical site.
Wound contamination from operating room personnel	Transition of microorganisms from the surgical personnel's shoes, mouths, or body can contaminate surgical wounds. ¹⁴ Microorganisms from the hands of health care workers in the operating room can move onto the patient and operating field if personnel do not perform appropriate handwashing or gloving. ^{14,46,47}
Wound contamination from surgical instruments	Sterilization eliminates all microorganisms on the surfaces of surgical instruments. Using insufficiently sterilized tools can lead to pathogen transmission. ⁴⁸

may accompany up to 44% of prosthetic joint infections.^{53,54} The presence of a sinus tract or purulent drainage has a specificity of between 97% and 100% and a positive predictive value of

100%.⁵⁵ Joint stiffness has a reported sensitivity of 20.5% and specificity of 99% in patients with a hematogenous source of prosthetic joint infection.⁵⁶ Many of the aforementioned

Table 3. Surgical Site Infection Prevention Strategies From Prospective Studies

Intervention	Type of studies	Absolute or median value	RR or OR (P value)
Preoperative			
Do not remove hair at the surgical site unless the presence of hair will affect the procedure ^a	Meta-analysis of 19 RCTs and 6 quasi-randomized trials ⁵⁹	- Razor vs clippers: 4.4% (84 of 1889) vs 2.5% (46 of 1834) - Razor vs depilatory cream: 7.8% (68 of 868) vs 3.6% (26 of 725) - Razor vs none: 4.2% (34 of 819) vs 2.1% (19 of 887)	- RR, 1.64 (.005) - RR, 2.28 (.02) - RR, 1.82 (.03)
Decolonize surgical patients with intranasal antistaphylococcal agent and antistaphylococcal skin antiseptic for high-risk procedures (eg, cardiothoracic, orthopedic) ^b	Meta-analysis of 5 RCTs and 12 observational studies ⁶⁰	Decolonization vs none: 0.8% (52 of 19 940) vs 2.0% (253 of 12 790)	RR, 0.41 (<.001)
Antimicrobial prophylaxis within 1 h of incision with weight-based antimicrobial agents selected based on most common pathogens for specific procedure ^{61,62c}	Cohort ⁶¹	- Administration within 30 min before incision vs 30-60 min before incision: 1.6% (22 of 1339) vs 2.4% (38 of 1558) - Administration within 30 min before incision vs after incision: 1.6% (22 of 1339) vs 5.2% (9 of 174)	- OR, 0.67 (.13) - OR, 3.27 (.003)
Use a checklist based on the World Health Organization 19-item surgical checklist to ensure adherence to best practices ^{63,64}	Multicenter, quasi-experimental study ⁶⁵	Without vs with checklist: 6.2% vs 3.4%	RR, 0.55 (<.001)
Intraoperative			
Using chlorhexidine gluconate and alcohol-containing skin preparatory agent in combination ^d	Meta-analysis of 4 RCTs ⁶⁶	Chlorhexidine gluconate + alcohol vs povidone iodine + alcohol: 4.0% (54 of 1337) vs 6.5% (86 of 1326)	RR, 0.62 (.005)
Maintain normothermia during the surgical procedure	Systematic review of 3 RCTs ⁶⁷	Hypothermia vs normothermia: 4.7% (14 of 299) vs 13% (37 of 290)	RR, 3.67 (.008)
Postoperative			
Maintain and monitor blood glucose levels regardless of diabetes status Maintain blood glucose values between 110 and 150 mg/dL	Meta-analysis of 15 RCTs ⁶⁸	Glycemic control (<150 mg/dL) vs conventional control (>150 mg/dL): 9.4% (231 of 2464) vs 16% (392 of 2488)	RR, 0.59 (<.001)
Application of incisional negative pressure wound dressings	Meta-analysis of 23 RCTs ⁶⁹	Incisional negative pressure wound therapy vs standard dressings: 9.7% (124 of 1279) vs 15% (191 of 1268)	RR, 0.67 (<.001)

Abbreviations: OR, odds ratio; RCT, randomized clinical trial; RR, relative risk.

SI conversion factor: To convert glucose from mg/dL to mmol/L, multiply by 0.0555.

^a If hair removal is necessary, remove outside of the operating room. For male genitalia, 1 RCT suggested that preoperative hair removal on scrotal skin using a razor as opposed to clippers resulted in less skin trauma without an increased risk in surgical site infection.⁷⁰

^b Antistaphylococcal skin antiseptic agents include chlorhexidine gluconate baths or wipes or dilute bleach baths.

^c Two hours are allowed for vancomycin and fluoroquinolones; redose antimicrobials for procedures with excessive blood loss and lengthy surgeries.

^d Alcohol containing skin preparation products are contraindicated for some procedures (eg, mucosa, cornea, or ear).

presentations may overlap with noninfectious conditions, such as a hematoma, seroma, or stitch abscess at points of suture penetration.

Classification of Surgical Site Infection

Despite variable presentations of surgical site infections, the US Centers for Disease Control and Prevention (CDC) National Healthcare Safety Network (NHSN) and the National Surgical Quality Improvement Program (NSQIP) provide specific surgical site infection definitions for surveillance and epidemiological purposes.^{57,58} Surveillance consists of systematic monitoring of patients following surgery to detect variance in surgical site infection rates and to develop quality improvement initiatives to lower infection rates. The goal of these definitions is to be simple and objective but flexible enough to encompass clinically relevant

infections. Both NHSN and NSQIP categorize surgical site infections into 3 groups: superficial-incisional (involving the skin or subcutaneous tissue layers of the incision), deep-incisional (involving muscle or connective tissue layers of the incision), and organs/spaces deep to the incision. Surveillance for surgical site infections continues for 30 days for most procedures and 90 days for specific procedures involving implanted materials. The NHSN collects data on all NHSN-eligible procedures, and NSQIP analyzes a subsample of 20% of cases for analysis via an 8-day systematic sampling cycle.

Prevention

Preoperative Period

A recent meta-analysis including 19 randomized and 6 quasi-randomized trials involving 8919 patients evaluated various ap-

proaches to preoperative hair removal for reducing surgical site infection (Table 3).⁵⁹ Across 7 randomized clinical trials (RCTs), hair removal with a razor was associated with a higher rate of surgical site infection: 4.4% (84 of 1889) patients whose hair was removed with a razor experienced an infection vs 2.5% (46 of 1834) whose hair was removed with clippers experienced an infection (relative risk [RR], 1.64 [95% CI, 1.16-2.33], $P = .005$). Across 9 RCTs, hair removal with a razor was associated with a higher rate of surgical site infection: 7.8% (68 of 868) patients vs 3.6% (26 of 725) patients whose hair was removed with a depilatory cream (RR, 2.28 [95% CI, 1.12-4.65]; $P = .02$). Seven RCTs demonstrated that removing hair with a razor was associated with an increased risk of surgical site infection: 4.2% (34 of 819) patients vs 2.1% (19 of 887) patients whose hair was not removed at all (RR, 1.82 [95% CI, 1.05-3.14]; $P = .03$).⁵⁹ Three RCTs reported that hair removal with clippers did not increase the risk of surgical site infection: 5.7% (49 of 863) patients vs 6.0% (52 of 870) patients whose hair was not removed (RR, 0.95 [95% CI, 0.65-1.39]; $P = .80$). If hair removal is necessary, it should be removed in the preoperative holding area and not in the operating room.

One method used to reduce surgical site infections is decolonization, in which patients are treated with an intranasal antimicrobial, skin antiseptic agent, or both to eliminate or temporarily reduce *S aureus* colonization prior to surgery. Evidence to support this recommendation is strongest for high-risk surgical procedures such as cardiothoracic surgeries and prosthetic joint replacement. This process typically includes an intranasal treatment with an antistaphylococcal agent (eg, mupirocin ointment or povidone iodine) and/or application of an antistaphylococcal skin antiseptic agent (eg, chlorhexidine gluconate solution or wipes) for 5 days. However, the precise timing, agent, and frequency of application are unclear because trials addressing this issue have used different strategies. The decolonization strategy should be completed as close to the surgical procedure as possible. A meta-analysis that included 5 RCTs and 12 observational studies showed that nasal decolonization was associated with lower rates of surgical site infections caused by gram-positive bacteria than no decolonization: 0.8% (152 of 19 940) vs 2.0% (253 of 12 790; RR, 0.41 [95% CI, 0.30-0.55]; $P < .001$).⁶⁰

This association persisted among the 11 studies in which patients were decolonized regardless of *S aureus* colonization status (RR, 0.40; 95% CI, 0.29-0.55) and among the 6 studies in which nasal decolonization was combined with skin antisepsis (RR, 0.29; 95% CI, 0.19-0.44, primary data not provided).⁶⁰ In contrast, other trials that included a more heterogeneous group of surgeries did not find a difference in surgical site infection incidence with decolonization.⁷¹ For example, a prospective cohort study that included 8 surgical categories (abdominal, orthopedic, urological, neurological, cardiovascular, thoracic, and plastic surgery and solid organ transplant) found that decolonization strategies did not reduce MRSA surgical site infections.⁷² The authors identified 60 MRSA infections (0.55%) among 10 910 procedures in the control group compared with 70 MRSA infections (0.65%) among 10 844 procedures during the intervention period ($P = .29$). As a result, decolonization is typically focused on orthopedic, cardiothoracic, or high-risk procedures such as spine and brain surgeries.

The intervention requires a significant amount of coordination to perform the appropriate test prior to surgery, have the result reviewed, and ensure the appropriate decolonization

approach was applied. Given the number of steps required, some hospitals perform decolonization on all patients undergoing high-risk surgical procedures, an approach that may ultimately be cost-effective (estimated \$153 per person) based on modeling studies.⁷³ In contrast, widespread use of antistaphylococcal antibiotics such as mupirocin may ultimately increase rates of resistant *S aureus* infections.^{74,75}

Conducting RCTs for surgical site infection prevention is challenging given the relatively low incidence of the outcome of interest. Thus, additional prevention strategies in the preoperative setting exist, but lack high-quality evidence. As a result, these interventions are predicated on expert opinion and results from retrospective cohort studies. For example, in contrast to postoperative glucose control, no RCTs have found a clear association between a specific hemoglobin A_{1c} cutoff and surgical site infections.

The administration of antibiotic prophylaxis is recommended in all surgical site infection prevention guidelines, despite the absence of RCTs.^{14,17,76,77} One multicenter cohort study involving 4186 patients found that risk of infection increased as the time from antibiotic infusion to incision increased, although the trend was not statistically significant: administration within 30 minutes prior to incision was associated with a risk of 1.6% (22 of 1339) vs 2.4% (38 of 1558) with administration of antibiotic between 31 and 60 minutes before surgery ($P = .13$).⁶¹ In the absence of trial data, guideline consensus is that antibiotics should be given within 60 minutes of the incision to maximize tissue concentration of the antibiotic. Additional recommendations include dosing antibiotics according to the patient's weight to ensure that adequate tissue concentrations are achieved and administering subsequent doses of antibiotics for lengthy procedures if excessive bleeding occurs. For example, cefazolin, the most commonly used agent for antimicrobial prophylaxis, should be redosed every 4 hours until completion of the procedure. These recommendations are mainly based on older cohort studies and evaluation of secondary outcomes (eg, tissue concentrations of antibiotics).⁶² Although the optimal duration of prophylactic antibiotics is not known, prolonged antimicrobial prophylaxis is increasingly associated with patient harm, such as acute kidney injury.⁷⁸ Authors of a systematic review of 28 randomized trials involving 9478 patients receiving either a single dose for prophylaxis or multiple doses concluded that additional doses did not reduce the risk of infection 6.2% (278 of 4499) vs 5.9% (261 of 4440; OR, 1.06 [95% CI, 0.89-1.25]).⁷⁹ Thus, guidelines recommend stopping antibiotic prophylactic antibiotics when the surgical wound is closed.

The WHO's surgical safety checklist is a 19-item list to improve adherence with best practice and decrease surgical site infection incidence. WHO developed this safety checklist to promote more consistent implementation of best practices. This 19-item checklist included surgical site infection (eg, antimicrobial prophylaxis) and non-surgical site infection components (eg, surgical time-out). A multicenter, quasi-experimental study of 8 sites and 3733 patients showed that the infection rate prior to the implementation of the checklist was 6.2% compared with 3.4% after implementation of the checklist (P value $< .001$ for the risk difference).⁶⁵ These results have been supported by subsequent multi- and single-center prospective studies.^{63,64} However, the exact mechanism of improvement is unclear and most likely multifactorial.

Intraoperative

Topical alcohol is highly bactericidal but does not have persistent activity when used as monotherapy for skin antisepsis (Table 3). Multiple guidelines recommend that surgical site antisepsis should be performed with a product that contains alcohol and another antiseptic agent (eg, chlorhexidine gluconate or povidone iodine).^{17,76,80} Products that combine alcohol and antiseptic agents are available in the US. Chlorhexidine gluconate plus alcohol appears to be superior to povidone iodine plus alcohol for the prevention of surgical site infections.⁸¹ In a meta-analysis of data from 4 RCTs involving 6916 women who had cesarean deliveries, the authors concluded that surgical site preparation with chlorhexidine gluconate plus alcohol was associated with lower rates of infection than preparation with povidone iodine plus alcohol: 4.0% (54 of 1337) vs 6.5% (86 of 1326; RR, 0.62 [95% CI, 0.45-0.87]; $P = .005$).⁶⁶ Similarly, a meta-analysis of 20 RCTs and 5 prospective, 4 retrospective, and 1 ambispective studies, including more than 29 000 participants found that skin preparation with chlorhexidine gluconate was associated with fewer surgical site infections than povidone iodine: 4.8% (725 of 15 263) vs 6.7% (925 of 13 743; RR, 0.65 [95% CI, 0.55-0.77]; $P < .001$).⁸²

Normothermia to keep core body temperatures from dropping during surgery is maintained by combinations of forced warm air, skin warming, and warmed intravenous fluids (Table 2). Targets for core temperatures vary: more than 35.5 °C and more than 36 °C. A systematic review of 3 RCTs examining active body surfacing warm systems for preventing complications of inadvertent perioperative hypothermia in adults found that using a forced air warming device was associated with lower rates of the risk of surgical site infection than no forced air warming: 4.7% (14 of 299) vs 13% (37 of 290; RR, 0.36 [95% CI, 0.20-0.66]; $P = .008$; Table 3).⁶⁷

Postoperative

Although there are no RCTs that have evaluated intensive glucose control to lower the preoperative average glucose (hemoglobin A_{1c}) vs usual care before surgery, postoperative hyperglycemia was associated with an increased risk of surgical site infections in patients with and without diabetes (Table 3).^{48,83,84} As a result, strategies to prevent hyperglycemia to prevent surgical site infection are recommended in all major guidelines. Most data to support this strategy are from RCTs involving patients with diabetes. In a meta-analysis of 15 RCTs comparing the use of tight glycemic control (<150 mg/dL; 8.32 mmol/L) with conventional control (>150 mg/dL), tight control was associated with lower rates of surgical site infection: 9.4% (231 of 2464) vs 16% (392 of 2488; RR, 0.59 [95% CI, 0.50-0.68]; $P < .001$).⁶⁸

Incisional negative pressure wound therapy, defined as wound dressing systems that continuously or intermittently apply subatmospheric pressure to the system, can reduce the risk of surgical site infection by promoting reducing fluid accumulation in the wounds, thereby accelerating primary wound healing. Authors of a meta-analysis of 23 RCTs involving 2547 patients undergoing various surgical procedures (eg, abdominal, cesarean delivery, orthopedic, vascular) concluded that use of incisional negative pressure wound therapy for primary wound closure was associated with lower rates of surgical site infection than use of standard dressings: 9.7% (124 of 1279) vs 15% (191 of 1268; RR, 0.67 [95% CI, 0.53-0.85]; $P < .001$); however, the effect varied by procedure type.⁶⁹ The authors indi-

cated that they did not find evidence for substantial differences between the different types of surgery. Similarly, authors of a recent meta-analysis of 28 RCTs concluded that incisional negative pressure wound therapy was associated with lower rates of surgical site infection than standard dressing: 8.8% (194 of 2193) vs 14% (315 of 2205; RR, 0.61 [95% CI, 0.49-0.76]; $P < .001$).⁸⁵ The authors specified that when stratified by surgical discipline, the greatest benefits for surgical site infection reduction occurred in vascular surgery (RR, 0.45; 95% CI, 0.32-0.65; $P < .001$) and cardiac surgery (RR, 0.17; 95% CI, 0.03-0.96; $P = .05$), whereas the intervention was not associated with statistically significant benefit for abdominal surgery (RR, 0.56; 95% CI, 0.30-1.03), obstetric surgery (RR, 0.73; 95% CI, 0.44-1.20), orthopedic or trauma-derived surgery (RR, 0.68; 95% CI, 0.43-1.08), and plastic surgery (RR, 0.82; 95% CI, 0.26-2.63). The broader CIs for these later 4 subgroups suggest the possibility that they were underpowered to find a significant difference.

Hospital-Wide Surveillance

As one of the original surgical site infection prevention investigations, data from the Study on the Efficacy of Nosocomial Infection Control (SENIC)⁸⁶ supported the use of routine surveillance and feedback to reduce infections. The multicenter, 1985 SENIC study, evaluated infection prevention practices and found that the use of standardized surgical site infection surveillance by trained infection prevention personnel and routine feedback to surgeons was associated with an estimated reduction in infections in US hospitals from 586 000 to 510 000 compared with when no surveillance and feedback were given. Current recommendations advise health care institutions to identify high-volume, high-risk procedures and implement a system for collecting and storing data. Periodic reports should be prepared and given to key stakeholders to provide feedback on infection rates. Surveillance and feedback, along with several other quality improvement strategies (eg, education of surgeons, surgical staff, and patients) are endorsed by all surgical site infection prevention guidelines.^{14,17,77,80}

Limitations

This review has several limitations. First, this review focused on prevention of surgical site infection following general, commonly performed surgical procedures. Second, not all recommendations in previously published guidelines were summarized herein given the lack of available RCT data. Third, some interventions had been studied in only a small number of RCTs. Fourth, in some cases, the only available studies were older. Fifth, quality of included literature was not assessed. Sixth, some relevant studies may have been missed.

Conclusions

Surgical site infections affect approximately 0.5% to 3% of patients undergoing surgery and are associated with longer hospital stays than patients with no surgical site infections. Avoiding razors for hair removal, maintaining normothermia, use of chlorhexidine gluconate plus alcohol-based skin preparation agents, decolonization with intranasal antistaphylococcal agents and antistaphylococcal skin antiseptics for high-risk procedures, controlling for perioperative glucose concentrations, and using negative pressure wound therapy can reduce the rate of surgical site infections.

ARTICLE INFORMATION

Accepted for Publication: December 13, 2022.

Conflict of Interest Disclosures: Dr Seidelman reported receiving royalties from UpToDate for its Pelvic Osteomyelitis page and providing expert testimony for legal case related to musculoskeletal infections. Dr Mantyh reported receiving personal fees from Becton Dickinson for advising outside the submitted work. Dr Anderson reported receiving grants from the CDC and royalties from UpToDate outside the submitted work; and owning Infection Control Education for Major Sports, LLC.

Submissions: We encourage authors to submit papers for consideration as a Review. Please contact Mary McGrae McDermott, MD, at mdm608@northwestern.edu.

REFERENCES

1. Surgical site infection event (SSI). National Healthcare Safety Network. Accessed April 26, 2022. <https://www.cdc.gov/nhsn/pdfs/pscmanual/9pscscurrent.pdf>
2. Karlens OE, Borgen P, Bragnes B, et al. Rifampin combination therapy in staphylococcal prosthetic joint infections: a randomized controlled trial. *J Orthop Surg Res*. 2020;15(1):365. doi:10.1186/s13018-020-01877-2
3. Dencker EE, Bonde A, Troelsen A, Varadarajan KM, Sillesen M. Postoperative complications: an observational study of trends in the United States from 2012 to 2018. *BMC Surg*. 2021;21(1):393. doi:10.1186/s12893-021-01392-z
4. Seidelman JL. Surgical site infection (SSI) trends in community hospitals from 2013 to 2018. *Infect Control Hosp Epidemiol*. Published online July 18, 2022. doi:10.1017/ice.2022.135
5. Gantz O, Zagadailov P, Merchant AM. The cost of surgical site infections after colorectal surgery in the United States from 2001 to 2012: a longitudinal analysis. *Am Surg*. 2019;85(2):142-149. doi:10.1177/000313481908500219
6. Magill SS, Edwards JR, Bannam W, et al; Emerging Infections Program Healthcare-Associated Infections and Antimicrobial Use Prevalence Survey Team. Multistate point-prevalence survey of health care-associated infections. *N Engl J Med*. 2014;370(13):1198-1208. doi:10.1056/NEJMoal306801
7. Zimlichman E, Henderson D, Tamir O, et al. Health care-associated infections: a meta-analysis of costs and financial impact on the US health care system. *JAMA Intern Med*. 2013;173(22):2039-2046. doi:10.1001/jamainternmed.2013.9763
8. O'Hara LM, Thom KA, Preas MA. Update to the Centers for Disease Control and Prevention and the Healthcare Infection Control Practices Advisory Committee Guideline for the prevention of surgical site infection (2017): a summary, review, and strategies for implementation. *Am J Infect Control*. 2018;46(6):602-609. doi:10.1016/j.ajic.2018.01.018
9. Scott RD. *The Direct Medical Costs of Healthcare-Associated Infections in US Hospitals and the Benefits of Prevention*. Centers for Disease Control and Prevention; March 2009. Publication C5200891-A. Accessed April 28, 2022. https://www.cdc.gov/hai/pdfs/hai/scott_costpaper.pdf
10. Anderson DJ, Kaye KS, Chen LF, et al. Clinical and financial outcomes due to methicillin resistant *Staphylococcus aureus* surgical site infection: a multi-center matched outcomes study. *PLoS One*. 2009;4(12):e8305. doi:10.1371/journal.pone.0008305
11. Ming DY, Chen LF, Miller BA, Anderson DJ. The impact of depth of infection and postdischarge surveillance on rate of surgical-site infections in a network of community hospitals. *Infect Control Hosp Epidemiol*. 2012;33(3):276-282. doi:10.1086/664053
12. Wenzel RP. Surgical site infections and the microbiome: an updated perspective. *Infect Control Hosp Epidemiol*. 2019;40(5):590-596. doi:10.1017/ice.2018.363
13. Seidelman JL, Baker AW, Lewis SS, Advani SD, Smith B, Anderson D; Duke Infection Control Outreach Network Surveillance Team. Surgical site infection trends in community hospitals from 2013 to 2018. *Infect Control Hosp Epidemiol*. Published online July 2022. doi:10.1017/ice.2022.135
14. Leaper DJ, Edmiston CE. World Health Organization: global guidelines for the prevention of surgical site infection. *J Hosp Infect*. 2017;95(2):135-136. doi:10.1016/j.jhin.2016.12.016
15. Seidelman JL, Ge M, Baker AW, et al. Colon surgical-site infections and the impact of "present at the time of surgery (PATOS)" in a large network of community hospitals. *Infect Control Hosp Epidemiol*. Published online September 22 2022. doi:10.1017/ice.2022.236
16. Zhang Y, Zheng QJ, Wang S, et al. Diabetes mellitus is associated with increased risk of surgical site infections: a meta-analysis of prospective cohort studies. *Am J Infect Control*. 2015;43(8):810-815. doi:10.1016/j.ajic.2015.04.003
17. Ban KA, Minei JP, Laronga C, et al. American College of Surgeons and Surgical Infection Society: surgical site infection guidelines, 2016 update. *J Am Coll Surg*. 2017;224(1):59-74. doi:10.1016/j.jamcollsurg.2016.10.029
18. Filsoufi F, Rahmani PB, Castillo JG, Pinney S, Broumand SR, Adams DH. Incidence, treatment strategies and outcome of deep sternal wound infection after orthotopic heart transplantation. *J Heart Lung Transplant*. 2007;26(11):1084-1090. doi:10.1016/j.healun.2007.07.036
19. Payne WG, Naidu DK, Wheeler CK, et al. Wound healing in patients with cancer. *Eplasty*. 2008;8:e9.
20. Hennessey DB, Burke JP, Ni-Dhonochu T, Shields C, Winter DC, Mealy K. Preoperative hypoalbuminemia is an independent risk factor for the development of surgical site infection following gastrointestinal surgery: a multi-institutional study. *Ann Surg*. 2010;252(2):325-329. doi:10.1097/SLA.0b013e3181e9819a
21. Meijis AP, Koek MBG, Vos MC, Geerlings SE, Vogely HC, de Greeff SC. The effect of body mass index on the risk of surgical site infection. *Infect Control Hosp Epidemiol*. 2019;40(9):991-996. doi:10.1017/ice.2019.165
22. Harrington G, Russo P, Spelman D, et al. Surgical-site infection rates and risk factor analysis in coronary artery bypass graft surgery. *Infect Control Hosp Epidemiol*. 2004;25(6):472-476. doi:10.1086/502424
23. Yuan K, Chen HL. Obesity and surgical site infections risk in orthopedics: a meta-analysis. *Int J Surg*. 2013;11(5):383-388. doi:10.1016/j.ijsu.2013.02.018
24. Mangram AJ, Horan TC, Pearson ML, Silver LC, Jarvis WR; Hospital Infection Control Practices Advisory Committee. Guideline for prevention of surgical site infection, 1999. *Infect Control Hosp Epidemiol*. 1999;20(4):250-278. doi:10.1086/501620
25. Wukich DK, McMillen RL, Lowery NJ, Frykberg RG. Surgical site infections after foot and ankle surgery: a comparison of patients with and without diabetes. *Diabetes Care*. 2011;34(10):2211-2213. doi:10.2337/dc11-0846
26. Kaye KS, Anderson DJ, Sloane R, et al. The effect of surgical site infection on older operative patients. *J Am Geriatr Soc*. 2009;57(1):46-54. doi:10.1111/j.1532-5415.2008.02053.x
27. Roig-Rosello E, Rousselle P. The human epidermal basement membrane: a shaped and cell instructive platform that aging slowly alters. *Biomolecules*. 2020;10(12):1607. doi:10.3390/biom10121607
28. Faraday N, Rock P, Lin EE, et al. Past history of skin infection and risk of surgical site infection after elective surgery. *Ann Surg*. 2013;257(1):150-154. doi:10.1097/SLA.0b013e3182588abf
29. Edmiston CE Jr, Seabrook GR, Cambria RA, et al. Molecular epidemiology of microbial contamination in the operating room environment: is there a risk for infection? *Surgery*. 2005;138(4):573-579. doi:10.1016/j.surg.2005.06.045
30. Lidwell OM, Lowbury EJ, Whyte W, Blowers R, Stanley SJ, Lowe D. Airborne contamination of wounds in joint replacement operations: the relationship to sepsis rates. *J Hosp Infect*. 1983;4(2):111-131. doi:10.1016/0195-6701(83)90041-5
31. Wang Z, Anderson FA Jr, Ward M, Bhattacharyya T. Surgical site infections and other postoperative complications following prophylactic anticoagulation in total joint arthroplasty. *PLoS One*. 2014;9(4):e91755. doi:10.1371/journal.pone.0091755
32. Rohde JM, Dimcheff DE, Blumberg N, et al. Health care-associated infection after red blood cell transfusion: a systematic review and meta-analysis. *JAMA*. 2014;311(13):1317-1326. doi:10.1001/jama.2014.2726
33. Wetterslev J, Meyhoff CS, Jørgensen LN, Gluud C, Lindschou J, Rasmussen LS. The effects of high perioperative inspiratory oxygen fraction for adult surgical patients. *Cochrane Database Syst Rev*. 2015;(6):CD008884. doi:10.1002/14651858.CD008884.pub2
34. Hopf HW, Holm J. Hyperoxia and infection. *Best Pract Res Clin Anaesthesiol*. 2008;22(3):553-569. doi:10.1016/j.bpa.2008.06.001
35. Berard F, Gandon J. Postoperative wound infections: the influence of ultraviolet irradiation of the operating room and of various other factors. *Ann Surg*. 1964;160(suppl 2):1-192.
36. Elek SD, Conen PE. The virulence of *Staphylococcus pyogenes* for man: a study of the problems of wound infection. *Br J Exp Pathol*. 1957;38(6):573-586.

37. Korol E, Johnston K, Waser N, et al. A systematic review of risk factors associated with surgical site infections among surgical patients. *PLoS One*. 2013;8(12):e83743. doi:10.1371/journal.pone.0083743
38. Kurz A, Sessler DI, Lenhardt R; Study of Wound Infection and Temperature Group. Perioperative normothermia to reduce the incidence of surgical-wound infection and shorten hospitalization. *N Engl J Med*. 1996;334(19):1209-1215. doi:10.1056/NEJM199605093341901
39. Latham R, Lancaster AD, Covington JF, Pirolo JS, Thomas CS Jr. The association of diabetes and glucose control with surgical-site infections among cardiothoracic surgery patients. *Infect Control Hosp Epidemiol*. 2001;22(10):607-612. doi:10.1086/501830
40. Mangram AJ, Horan TC, Pearson ML, Silver LC, Jarvis WR, Committee HICPA; Hospital Infection Control Practices Advisory Committee. Guideline for prevention of surgical site infection, 1999. *Infect Control Hosp Epidemiol*. 1999;20(4):250-278. doi:10.1086/501620
41. Dumville JC, Gray TA, Walter CJ, et al. Dressings for the prevention of surgical site infection. *Cochrane Database Syst Rev*. 2016;12(12):CD003091. doi:10.1002/14651858.CD003091.pub4
42. Manian FA. The role of postoperative factors in surgical site infections: time to take notice. *Clin Infect Dis*. 2014;59(9):1272-1276. doi:10.1093/cid/ciu552
43. McLaws ML, Murphy C, Whitby M; Hospital Infection Standardised Surveillance. Standardising surveillance of nosocomial infections: the HISS program. *J Qual Clin Pract*. 2000;20(1):6-11. doi:10.1046/j.1440-1762.2000.00347.x
44. Tammelin A, Hambraeus A, Ståhle E. Source and route of methicillin-resistant *Staphylococcus epidermidis* transmitted to the surgical wound during cardio-thoracic surgery: possibility of preventing wound contamination by use of special scrub suits. *J Hosp Infect*. 2001;47(4):266-276. doi:10.1053/jhin.2000.0914
45. Lee KY, Coleman K, Paech D, Norris S, Tan JT. The epidemiology and cost of surgical site infections in Korea: a systematic review. *J Korean Surg Soc*. 2011;81(5):295-307. doi:10.4174/jkss.2011.81.5.295
46. Fagernes M, Lingaas E. Factors interfering with the microflora on hands: a regression analysis of samples from 465 healthcare workers. *J Adv Nurs*. 2011;67(2):297-307. doi:10.1111/j.1365-2648.2010.05462.x
47. Loftus RW, Muffy MK, Brown JR, et al. Hand contamination of anesthesia providers is an important risk factor for intraoperative bacterial transmission. *Anesth Analg*. 2011;112(1):98-105. doi:10.1213/ANE.0b013e3181e7ce18
48. Kotagal M, Symons RG, Hirsch IB, et al; SCOAP-CERTAIN Collaborative. Perioperative hyperglycemia and risk of adverse events among patients with and without diabetes. *Ann Surg*. 2015;261(1):97-103. doi:10.1097/SLA.0000000000000688
49. Aghdassi SJS, Schröder C, Gastmeier P. Gender-related risk factors for surgical site infections: results from 10 years of surveillance in Germany. *Antimicrob Resist Infect Control*. 2019;8:95. doi:10.1186/s13756-019-0547-x
50. Anderson DJ, Arduino JM, Reed SD, et al. Variation in the type and frequency of postoperative invasive *Staphylococcus aureus* infections according to type of surgical procedure. *Infect Control Hosp Epidemiol*. 2010;31(7):701-709. doi:10.1086/653205
51. Ghosh S, Charity RM, Haidar SG, Singh BK. Pyrexia following total knee replacement. *Knee*. 2006;13(4):324-327. doi:10.1016/j.knee.2006.05.001
52. Zajonc D, Wuthe L, Tiepolt S, et al. Diagnostic work-up strategy for periprosthetic joint infections after total hip and knee arthroplasty: a 12-year experience on 320 consecutive cases. *Patient Saf Surg*. 2015;9:20. doi:10.1186/s13037-015-0071-8
53. Berbari EF, Hanssen AD, Duffy MC, et al. Risk factors for prosthetic joint infection: case-control study. *Clin Infect Dis*. 1998;27(5):1247-1254. doi:10.1086/514991
54. Berbari EF, Osmon DR, Carr A, et al. Dental procedures as risk factors for prosthetic hip or knee infection: a hospital-based prospective case-control study. *Clin Infect Dis*. 2010;50(1):8-16. doi:10.1086/648676
55. Portillo ME, Salvadó M, Sorli L, et al. Multiplex PCR of sonication fluid accurately differentiates between prosthetic joint infection and aseptic failure. *J Infect*. 2012;65(6):541-548. doi:10.1016/j.jinf.2012.08.018
56. Tande AJ, Palraj BR, Osmon DR, et al. Clinical presentation, risk factors, and outcomes of hematogenous prosthetic joint infection in patients with staphylococcus aureus bacteremia. *Am J Med*. 2016;129(2):221.E11-221.E20. doi:10.1016/j.amjmed.2015.09.006
57. Wilson BM, Bessesen MT, Doros G, et al. Adjunctive rifampin therapy for diabetic foot osteomyelitis in the Veterans Health Administration. *JAMA Netw Open*. 2019;2(11):e1916003. doi:10.1001/jamanetworkopen.2019.16003
58. Zhu C, Leach JR, Wang Y, Gasper W, Saloner D, Hope MD. Intraluminal thrombus predicts rapid growth of abdominal aortic aneurysms. *Radiology*. 2020;294(3):707-713. doi:10.1148/radiol.2020191723
59. Tanner J, Melen K. Preoperative hair removal to reduce surgical site infection. *Cochrane Database Syst Rev*. 2021;8(8):CD004122. doi:10.1002/14651858.CD004122.pub5
60. Schweizer M, Perencevich E, McDanel J, et al. Effectiveness of a bundled intervention of decolonization and prophylaxis to decrease Gram positive surgical site infections after cardiac or orthopedic surgery: systematic review and meta-analysis. *BMJ*. 2013;346:f2743. doi:10.1136/bmj.f2743
61. Steinberg JP, Braun BI, Hellinger WC, et al; Trial to Reduce Antimicrobial Prophylaxis Errors (TRAPE) Study Group. Timing of antimicrobial prophylaxis and the risk of surgical site infections: results from the Trial to Reduce Antimicrobial Prophylaxis Errors. *Ann Surg*. 2009;250(1):10-16. doi:10.1097/SLA.0b013e3181ad5fca
62. Bratzler DW, Dellinger EP, Olsen KM, et al; American Society of Health-System Pharmacists; Infectious Disease Society of America; Surgical Infection Society; Society for Healthcare Epidemiology of America. Clinical practice guidelines for antimicrobial prophylaxis in surgery. *Am J Health Syst Pharm*. 2013;70(3):195-283. doi:10.2146/ajhp120568
63. van Klei WA, Hoff RG, van Aarnhem EE, et al. Effects of the introduction of the WHO "Surgical Safety Checklist" on in-hospital mortality: a cohort study. *Ann Surg*. 2012;255(1):44-49. doi:10.1097/SLA.0b013e31823779ae
64. Weiser TG, Haynes AB, Dziekan G, Berry WR, Lipsitz SR, Gawande AA; Safe Surgery Saves Lives Investigators and Study Group. Effect of a 19-item surgical safety checklist during urgent operations in a global patient population. *Ann Surg*. 2010;251(5):976-980. doi:10.1097/SLA.0b013e3181d970e3
65. Haynes AB, Weiser TG, Berry WR, et al; Safe Surgery Saves Lives Study Group. A surgical safety checklist to reduce morbidity and mortality in a global population. *N Engl J Med*. 2009;360(5):491-499. doi:10.1056/NEJMsa0801019
66. Hadiati DR, Hakimi M, Nurdianti DS, Masuzawa Y, da Silva Lopes K, Ota E. Skin preparation for preventing infection following caesarean section. *Cochrane Database Syst Rev*. 2020;6(6):CD007462. doi:10.1002/14651858.CD007462.pub5
67. Madrid E, Urrutia G, Roqué i Figuls M, et al. Active body surface warming systems for preventing complications caused by inadvertent perioperative hypothermia in adults. *Cochrane Database Syst Rev*. 2016;4(4):CD009016. doi:10.1002/14651858.CD009016.pub2
68. Wang YY, Hu SF, Ying HM, et al. Postoperative tight glycemic control significantly reduces postoperative infection rates in patients undergoing surgery: a meta-analysis. *BMC Endocr Disord*. 2018;18(1):42. doi:10.1186/s12902-018-0268-9
69. Norman G, Shi C, Goh EL, et al. Negative pressure wound therapy for surgical wounds healing by primary closure. *Cochrane Database Syst Rev*. 2022;4(4):CD009261. doi:10.1002/14651858.CD009261.pub7
70. Grober ED, Domes T, Fanipour M, Copp JE. Preoperative hair removal on the male genitalia: clippers vs razors. *J Sex Med*. 2013;10(2):589-594. doi:10.1111/j.1743-6109.2012.02904.x
71. Perl TM, Cullen JJ, Wenzel RP, et al; Mupirocin and the Risk of *Staphylococcus Aureus* Study Team. Intranasal mupirocin to prevent postoperative *Staphylococcus aureus* infections. *N Engl J Med*. 2002;346(24):1871-1877. doi:10.1056/NEJMoa003069
72. Harbarth S, Fankhauser C, Schrenzel J, et al. Universal screening for methicillin-resistant *Staphylococcus aureus* at hospital admission and nosocomial infection in surgical patients. *JAMA*. 2008;299(10):1149-1157. doi:10.1001/jama.299.10.1149
73. Rennert-May E, Conly J, Smith S, et al. A cost-effectiveness analysis of mupirocin and chlorhexidine gluconate for *Staphylococcus aureus* decolonization prior to hip and knee arthroplasty in Alberta, Canada, compared to standard of care. *Antimicrob Resist Infect Control*. 2019;8:113. doi:10.1186/s13756-019-0568-5
74. Kline SE, Sanstead EC, Johnson JR, Kulasingam SL. Cost-effectiveness of pre-operative *Staphylococcus aureus* screening and decolonization. *Infect Control Hosp Epidemiol*. 2018;39(11):1340-1346. doi:10.1017/ice.2018.228

75. Stambough JB, Nam D, Warren DK, et al. Decreased hospital costs and surgical site infection incidence with a universal decolonization protocol in primary total joint arthroplasty. *J Arthroplasty*. 2017;32(3):728-734.e1. doi:10.1016/j.arth.2016.09.041
76. Berríos-Torres SI, Umscheid CA, Bratzler DW, et al; Healthcare Infection Control Practices Advisory Committee. Centers for Disease Control and Prevention guideline for the prevention of surgical site infection, 2017. *JAMA Surg*. 2017;152(8):784-791. doi:10.1001/jamasurg.2017.0904
77. Anderson DJ, Podgorny K, Berríos-Torres SI, et al. Strategies to prevent surgical site infections in acute care hospitals: 2014 update. *Infect Control Hosp Epidemiol*. 2014;35(6):605-627. doi:10.1086/676022
78. Branch-Elliman W, O'Brien W, Strymish J, Itani K, Wyatt C, Gupta K. Association of duration and type of surgical prophylaxis with antimicrobial-associated adverse events. *JAMA Surg*. 2019;154(7):590-598. doi:10.1001/jamasurg.2019.0569
79. McDonald M, Grabsch E, Marshall C, Forbes A. Single- versus multiple-dose antimicrobial prophylaxis for major surgery: a systematic review. *Aust N Z J Surg*. 1998;68(6):388-396. doi:10.1111/j.1445-2197.1998.tb04785.x
80. World Health Organization. *Global Guidelines for the Prevention of Surgical Site Infection*. World Health Organization; 2016.
81. Darouiche RO, Wall MJ Jr, Itani KM, et al. Chlorhexidine-alcohol versus povidone-iodine for surgical-site antisepsis. *N Engl J Med*. 2010;362(1):18-26. doi:10.1056/NEJMoa0810988
82. Chen S, Chen JW, Guo B, Xu CC. Preoperative antisepsis with chlorhexidine versus povidone-iodine for the prevention of surgical site infection: a systematic review and meta-analysis. *World J Surg*. 2020;44(5):1412-1424. doi:10.1007/s00268-020-05384-7
83. Kao LS, Phatak UR. Glycemic control and prevention of surgical site infection. *Surg Infect (Larchmt)*. 2013;14(5):437-444. doi:10.1089/sur.2013.008
84. Kiran RP, Turina M, Hammel J, Fazio V. The clinical significance of an elevated postoperative glucose value in nondiabetic patients after colorectal surgery: evidence for the need for tight glucose control? *Ann Surg*. 2013;258(4):599-604. doi:10.1097/SLA.0b013e3182a501e3
85. Zwanenburg PR, Tol BT, Obdeijn MC, Lapid O, Gans SL, Boermeester MA. Meta-analysis, meta-regression, and GRADE assessment of randomized and nonrandomized studies of incisional negative pressure wound therapy versus control dressings for the prevention of postoperative wound complications. *Ann Surg*. 2020;272(1):81-91. doi:10.1097/SLA.0000000000003644
86. Haley RW, Culver DH, White JW, et al. The efficacy of infection surveillance and control programs in preventing nosocomial infections in US hospitals. *Am J Epidemiol*. 1985;121(2):182-205. doi:10.1093/oxfordjournals.aje.a113990