

Ovarian Cancer

A Review

Giuseppe Caruso, MD; S. John Weroha, MD, PhD; William Cliby, MD

IMPORTANCE Ovarian cancer is the eighth most common cause of cancer and cancer death in women worldwide. In 2022, ovarian cancer was diagnosed in approximately 324 398 individuals, and 206 839 died of ovarian cancer worldwide. In 2025, it is estimated that 20 890 US women will be diagnosed with ovarian cancer and 12 730 patients will die of ovarian cancer.

OBSERVATIONS Approximately 90% of ovarian cancers are epithelial malignancies, of which 70% to 80% are high-grade serous ovarian cancers. Less common epithelial subtypes include endometrioid, clear cell, low-grade serous, mucinous, and carcinosarcoma. The median age at diagnosis of ovarian cancer is 63 years. Risk factors include older age, family history of breast or ovarian cancer, endometriosis, and nulliparity. Hereditary factors are associated with 25% of cases, predominantly linked to *BRCA1/2* gene variants. At diagnosis, approximately 95% of patients experience nonspecific symptoms, such as abdominal pain, bloating, and urinary urgency and frequency, and about 80% have advanced-stage disease (stage III-IV), including extrapelvic disease, ascites, and abdominal masses. Diagnostic and staging evaluation includes pelvic ultrasound; computed tomography of the chest, abdomen, and pelvis; and serum tumor markers such as carbohydrate antigen 125, carbohydrate antigen 19-9, and carcinoembryonic antigen. First-line treatment for early-stage ovarian cancer, defined as limited to the ovary or fallopian tube (stage I) or confined to the pelvis (stage II), is surgery (hysterectomy, bilateral salpingo-oophorectomy, omentectomy, and lymphadenectomy), followed by adjuvant chemotherapy (carboplatin and paclitaxel). With treatment, early-stage ovarian cancer has a 5-year overall survival of 70% to 95%. Advanced-stage ovarian cancer may be treated with primary cytoreductive surgery (removal of all visible cancer in the abdominal cavity) and adjuvant chemotherapy (carboplatin and paclitaxel) or with neoadjuvant chemotherapy followed by cytoreductive surgery and adjuvant chemotherapy. Most patients with advanced-stage ovarian cancer receive maintenance therapy with bevacizumab (a monoclonal antibody that blocks angiogenesis) and/or poly-adenosine diphosphate ribose polymerase (PARP) inhibitors. With treatment, the 5-year overall survival rate for advanced-stage ovarian cancer is 10% to 40%. However, individuals with *BRCA*-related gene variants have a 5-year overall survival rate of approximately 70% with PARP inhibitor treatment. Despite an initial remission rate of 80%, approximately 75% of patients with advanced-stage disease have ovarian cancer relapse within 2 years.

CONCLUSIONS AND RELEVANCE Approximately 21 000 women are diagnosed with ovarian cancer annually in the US, and approximately 80% have advanced-stage ovarian cancer at diagnosis. First-line treatment of early-stage ovarian cancer is surgery and adjuvant platinum-based chemotherapy. Treatment of advanced-stage ovarian cancer includes cytoreductive surgery, platinum-based chemotherapy, and targeted maintenance therapies such as bevacizumab and/or PARP inhibitors.

JAMA. 2025;334(14):1278-1291. doi:10.1001/jama.2025.9495
Published online July 21, 2025.

- [+ Multimedia](#)
- [+ Supplemental content](#)
- [+ CME at \[jamacmelookup.com\]\(https://jamacmelookup.com\)](#)

Author Affiliations: Department of Obstetrics and Gynecology, Division of Gynecologic Oncology, Mayo Clinic, Rochester, Minnesota (Caruso, Cliby); Division of Gynecologic Oncology, European Institute of Oncology (IEO), IRCCS, Milan, Italy (Caruso); Department of Experimental Medicine, Sapienza University of Rome, Rome, Italy (Caruso); Department of Oncology, Mayo Clinic, Rochester, Minnesota (Weroha).

Corresponding Author: William Cliby, MD, Department of Obstetrics and Gynecology, Division of Gynecologic Surgery, Mayo Clinic, 200 First St SW, Rochester, MN 55905 (cliby.william@mayo.edu).

Among US women, the lifetime risk of developing ovarian cancer is 1 in 91, and death due to ovarian cancer occurs 1 in 143 women.¹ Approximately 80% of ovarian cancers are at an advanced stage at diagnosis.²⁻⁴ Earlier detection is difficult because there are no effective screening methods, and 95% of patients have nonspecific initial symptoms, such as abdominal pain, bloating, and urinary urgency or frequency.

Epithelial malignancies comprise 90% of ovarian cancers, with 70% to 80% being high-grade serous ovarian cancers.⁵ Early-stage ovarian cancer includes stage I (confined to the ovary or fallopian tube) and stage II (involving the uterus or other pelvic organs) (Figure 1). With surgery and adjuvant platinum-based chemotherapy, the 5-year overall survival of patients with early-stage ovarian cancer is 70% to 95%.^{2,4} Advanced-stage ovarian cancer, which comprises stage III (involving pelvic and para-aortic lymph nodes or extrapelvic peritoneum) and stage IV (metastases to the liver or spleen or outside the abdominal cavity), may be treated with a combination of cytoreductive surgery, platinum-based chemotherapy, and targeted therapies such as the antiangiogenic drug bevacizumab and/or poly-adenosine diphosphate ribose polymerase (PARP) inhibitors. Initial remission rates of advanced-stage ovarian cancer are 80%, but approximately 75% of patients with advanced-stage ovarian cancer relapse within 2 years, and 5-year overall survival is 10% to 40%.^{2,4} Approximately 50% of patients with advanced-stage ovarian cancer have heritable (germline) or tumor-acquired (somatic) *BRCA1/2* or other *BRCA*-related gene variants, and when treated with PARP inhibitors, have 5-year overall survival rates approaching 70%.⁶ This Review focuses on diagnosis and manage-

ment of high-grade serous ovarian cancer. Box 1 provides some common questions and answers about ovarian cancer.


Methods

A PubMed search of English-language articles describing randomized clinical trials (RCTs), meta-analyses, systematic reviews, and practice guidelines published between January 1, 1995, and May 3, 2025, identified 1733 RCTs, 1457 meta-analyses, 1770 systematic reviews, and 295 guidelines. Priority was given to the selection of high-quality RCTs and meta-analyses. A total of 123 articles were included, comprising 65 RCTs, 9 meta-analyses, 5 systematic reviews, 16 guidelines, 6 narrative reviews, 12 prospective observational studies, and 10 retrospective observational studies.

Epidemiology and Risk Factors

Ovarian cancer is the eighth most common cause of cancer and cancer death in women worldwide.⁷ Globally, in 2022, there were an estimated 324 398 new cases of ovarian cancer and 206 839 deaths, with incidence and mortality rates of 6.6 and 4.2 per 100 000 women-years, respectively.⁷ In the US, ovarian cancer incidence and mortality rates have declined by approximately 3% annually since 2004 due to increased use of oral contraceptives (preventing repetitive ovulation-related ovarian injury), prophylactic salpingo-oophorectomy in women with high-risk genetic variants such as

Figure 1. Origin and Stages of Ovarian Cancer



Box 1. Commonly Asked Questions About Ovarian Cancer**What Are Common Presenting Signs and Symptoms of Ovarian Cancer?**

Up to 95% of women with ovarian cancer initially experience nonspecific symptoms, such as abdominal pain, bloating, and urinary urgency. Signs and symptoms of advanced-stage ovarian cancer may include ascites, palpable abdominal masses, and/or dyspnea (caused by abdominal distension or pleural effusions).

Are There Effective Methods to Decrease Risk of Ovarian Cancer?

The US Preventive Services Task Force in 2018 recommended against ovarian cancer screening with transvaginal ultrasound and/or carbohydrate antigen 125 in both asymptomatic individuals without a high-risk hereditary cancer syndrome and high-risk individuals (eg, *BRCA1/2* gene variant carriers). Women with a family history of a *BRCA* gene variant or with a first-degree relative with ovarian cancer at any age or early-onset breast cancer (age <40 years) should undergo genetic testing. Bilateral salpingo-oophorectomy is recommended between ages 35 and 40 years for *BRCA1* carriers who have completed childbearing, between ages 40 and 45 years for *BRCA2* carriers, and between ages 45 and 50 years for women with Lynch syndrome.

What Is the Initial Treatment and Prognosis for Patients With Ovarian Cancer?

First-line treatment for early-stage ovarian cancer (stages I-II) is surgery followed by platinum-based chemotherapy, which is associated with 5-year overall survival rates of 70% to 95%. Advanced-stage ovarian cancer (stages III-IV) is treated with a combination of cytoreductive surgery, platinum-based chemotherapy, and targeted therapies (bevacizumab and/or PARP inhibitors) and has a 5-year survival rate of 10% to 40%. However, with PARP inhibitor treatment, patients with advanced-stage ovarian cancer and *BRCA*-related gene variants have a 5-year survival rate of approximately 70%.

BRCA1/2 (hereditary breast and ovarian cancer syndrome), and improved treatment options.¹ Globally, ovarian cancer incidence has decreased in other higher-income countries such as the UK, Austria, the Netherlands, and Norway but has increased in some lower-income regions such as Africa and some parts of Asia.^{7,8}

Ovarian cancer is predominantly of epithelial origin (90%). The World Health Organization in 2020 recognized 6 principal epithelial subtypes: high-grade serous (70%-80%), endometrioid (10%-20%), clear cell (5%-10%), low-grade serous (5%-10%), mucinous (3%-5%), and carcinosarcoma (1%-3%).⁵ Less commonly, ovarian cancer involves germ cell tumors (3%-5%) and sex cord stromal tumors (2%-5%).⁵

The median age at diagnosis of ovarian cancer is 63 years, although it is 10 years earlier in patients with germline *BRCA* gene variants.⁹ Approximately 25% of ovarian cancers are hereditary due to germline pathogenic variants in DNA damage repair genes: 90% in the *BRCA1* and *BRCA2* genes¹⁰ and 10% attributable to other gene variants, including the DNA mismatch repair genes (Lynch syndrome) and genes involved in the DNA double-strand breaks repair system, such as *CHEK2*, *RAD51*, *BRIP1*, and *PALB2*.¹¹ The estimated lifetime risk of developing ovarian cancer in patients with a *BRCA* gene variant is 20% to 70% and is 10% to 18% with a mismatch repair gene variant.^{12,13}

Endocrine and reproductive risk factors for ovarian cancer include endometriosis (hazard ratio, 4.20; 95% CI, 3.59-4.91),¹⁴ infertility (standardized incidence ratio, 2.0; 95% CI, 1.8-4.0), and postmenopausal estrogen therapy (relative risk, 1.31; 95% CI, 1.21-1.41).¹⁵ Possible protective factors are multiparity (odds ratio for each additional birth, 0.81; 95% CI, 0.75-0.85), breastfeeding (odds ratio, 0.72; 95% CI, 0.68-0.76), and oral contraceptive use (odds ratio, 0.66; 95% CI, 0.52-0.83).¹⁵

Pathophysiology and Molecular Biology

The pathogenesis of epithelial ovarian cancers is not clear, but gene expression profiling suggests that most high-grade serous ovarian cancers likely arise from precursors in the tubal fimbria and metastasize to the ovary and peritoneal cavity (Figure 1).¹⁶ About 50% of high-grade serous ovarian cancers have genetic variants (25% heritable/germline and 25% tumor-acquired/somatic) in homologous recombination repair genes resulting in an inability to repair DNA double-strand breaks, known as homologous recombination deficiency (HRD). *BRCA1* and *BRCA2* gene variants account for approximately half of HRD cases, found in 25% of all high-grade serous ovarian cancers.¹⁷ The remaining half of HRD cases are caused by other genes involved in the homologous recombination repair pathway, including *CHEK2*, *RAD51*, *BRIP1*, and *PALB2*.

Prevention**High-Risk Individuals**

Individuals at high risk of ovarian cancer include those with germline *BRCA1/2* gene variants and those with a family history of ovarian cancer at any age or early-onset breast cancer (age <40 years). The National Comprehensive Cancer Network (NCCN) guidelines recommend genetic testing for ovarian cancer susceptibility genes (*ATM*, *BRCA1*, *BRCA2*, *BRIP1*, *MLH1*, *MSH2*, *MSH6*, *EPCAM*, *PALB2*, *RAD51C*, and *RAD51D*) in individuals with a first- or second-degree relative with epithelial ovarian cancer at any age or with a personal or family cancer history suggestive of hereditary breast and ovarian cancer syndrome based on genetic counseling.¹⁰

For patients with known ovarian cancer susceptibility genes, prophylactic bilateral salpingo-oophorectomy is the most effective preventive measure for ovarian cancer.¹⁸ A meta-analysis of 3 prospective studies including 9192 women with *BRCA1/2* variants reported that bilateral salpingo-oophorectomy reduced the risk of ovarian cancer by 81% over 4 years (hazard ratio, 0.19; 95% CI, 0.13-0.27).¹⁹ The NCCN guidelines recommend bilateral salpingo-oophorectomy at age 35 to 40 years for *BRCA1* gene variant carriers who have completed childbearing, age 40 to 45 years for *BRCA2* carriers, and age 45 to 50 years for women with other pathogenic gene variants, such as for Lynch syndrome.¹⁰ Prior to prophylactic surgery, clinicians should discuss the potential effects of early menopause (eg, increased risk of vasomotor symptoms, osteoporosis, cardiovascular disease, cognitive decline, adverse effects on sexual function), and hormone therapy should be considered for premenopausal women with no personal breast cancer history.¹⁰

Oral contraceptives reduce ovarian cancer risk for high-risk younger individuals (eg, *BRCA* variant carriers) who do not undergo bilateral salpingo-oophorectomy. A meta-analysis of women with

BRCA1/2 variants using oral contraceptives reported a hazard ratio of 0.62 (95% CI, 0.52-0.74) for ovarian cancer based on 2 studies of 10 981 women and an odds ratio of 0.49 (95% CI, 0.38-0.63) from 8 studies including 10 390 women.²⁰ Similarly, another meta-analysis of 5 observational studies including 1503 participants with *BRCA1/2* pathogenic variants reported that oral contraceptive use was associated with a decreased risk of ovarian cancer (summary relative risk, 0.50; 95% CI, 0.33-0.75), and each additional 10 years of use was associated with a 36% reduced risk (summary relative risk, 0.64; 95% CI, 0.53-0.78).²¹

General Population

The American College of Obstetricians and Gynecologists and International Federation of Gynecology and Obstetrics guidelines recommend considering surgical and hormonal preventive approaches for the general population to reduce ovarian cancer risk.^{22,23} For postmenopausal women undergoing pelvic surgery for benign conditions (eg, hysterectomy for uterine fibroids or pelvic surgery for colon disease), bilateral salpingo-oophorectomy may be considered. For premenopausal women not interested in childbearing who are undergoing pelvic surgery, bilateral salpingectomy (removal of the fallopian tubes) while preserving the ovaries can avoid early menopause and its potential adverse effects on health.^{22,23}

A meta-analysis of 24 case-control and cohort studies reported a reduced incidence of ovarian cancer in oral contraceptive ever users compared with never users (odds ratio, 0.73; 95% CI, 0.66-0.81). Oral contraceptive use for more than 10 years was associated with lower risk (odds ratio, 0.43; 95% CI, 0.37-0.51).²⁴ Lifetime ovarian cancer reduction attributable to oral contraceptive use was estimated at 0.54% (number needed to treat, 185 for 5 years of oral contraceptive use).²⁴

Screening

The US Preventive Services Task Force in 2018 recommended against ovarian cancer screening in asymptomatic women without high-risk hereditary syndromes based on 2 large RCTs (PLCO and UKCTOCS; N = 271 103) that found no significant mortality reduction with use of annual transvaginal ultrasound and/or carbohydrate antigen 125 (CA125) testing vs no screening among average-risk, asymptomatic women (PLCO: relative risk, 1.18; 95% CI, 0.82-1.71; UKCTOCS: hazard ratio, 0.91; 95% CI, 0.76-1.09).²⁵ Similarly, large prospective studies of high-risk individuals (eg, those with *BRCA* gene variants) have shown that screening with transvaginal ultrasound and CA125 has a low positive predictive value (4.6%-10.8%) and does not reduce ovarian cancer mortality.²⁶⁻²⁹ Thus, screening for ovarian cancer is not recommended, even in high-risk individuals.

Clinical Presentation

Prior to ovarian cancer diagnosis, 95% of women report a gradual onset of nonspecific symptoms, such as abdominal pain, bloating, and urinary urgency or frequency.³⁰ A prospective case-control study of symptoms in women with a pelvic mass (84 benign and 44 ovarian cancer) compared with women visiting 2 primary care clinics (n = 1709) found that patients with ovarian cancer reported more fre-

quent bloating (70% vs 38%; $P < .001$), increased abdominal size (64% vs 19%; $P < .001$), urinary urgency/frequency (55% vs 32%; $P = .002$), abdominal pain (50% vs 30%; $P = .006$), and pelvic pain (41% vs 26%; $P = .02$).³¹ The combination of bloating, increased abdominal size, and urinary urgency/frequency was observed in 43% of patients with ovarian cancer vs 8% in patients without ovarian cancer (odds ratio, 9.4; 95% CI, 5.0-17.7).³¹ In a survey of 1725 US and Canadian women with ovarian cancer, 89% of patients with early-stage disease and 97% of patients with advanced-stage disease reported symptoms before diagnosis, with no significant difference in symptom type based on stage.³⁰ Persistent abdominal pain or bloating, ascites, a palpable abdominal or pelvic mass, weight loss, and dyspnea (caused by abdominal distension or pleural effusions) may indicate advanced-stage ovarian cancer. Rarely (<1% of patients), ovarian cancer presents with extra-abdominal symptoms caused by distant metastases (bone pain, seizures) or paraneoplastic syndromes (cerebellar degeneration, peripheral polyneuropathy).³²

Diagnosis

Diagnostic workup includes eliciting a personal and family history of cancer to determine if patients have a known genetic variant associated with ovarian cancer, such as *BRCA1/2*, and a physical examination, including pelvic and abdominal evaluations for palpable masses or ascites (Box 2). Pelvic ultrasound is highly sensitive (70%-93%) and specific (80%-98%) for identification of malignant adnexal masses.^{33,34} Computed tomography (CT) of the abdomen/pelvis is less sensitive (60%-90%) and specific (85%-94%) for diagnosis but has a sensitivity of 94% for detecting peritoneal disease and omental thickening, which are important findings for staging and planning.^{34,35} A detailed staging system for ovarian cancer is outlined in eTable 1 in the Supplement.³⁶ Compared with CT, abdominal magnetic resonance imaging (MRI) is more sensitive (92%-94%) and specific (85%-98%) for evaluating peritoneal carcinomatosis and mesenteric or bowel serosal involvement.^{34,35,37} If uncertainty remains about cancer resectability after CT or MRI imaging, diagnostic laparoscopy may be performed prior to laparotomy to determine resectability.³⁸ Total-body positron emission tomography has higher sensitivity (73%-75%) than CT (43%-55%) for detection of extra-abdominal lesions, such as distant lymph nodes (eg, mediastinal, inguinal, supraclavicular) and pulmonary metastases, potentially altering management to systemic therapy instead of abdominal debulking.^{39,40}

Serum Tumor Markers

Serum tumor markers, such as CA125, human epididymis protein 4, carbohydrate antigen 19-9 (CA19-9), and carcinoembryonic antigen (CEA), are not diagnostic of epithelial ovarian cancer but can help monitor treatment response and posttreatment surveillance. Carbohydrate antigen 125 is the most commonly elevated serum tumor marker,⁴¹ and based on 17 studies that included 2374 women with ovarian tumors, CA125 (≥ 35 U/mL) was elevated at diagnosis in approximately 50% of patients with early-stage ovarian cancer and in 75% to 92% with advanced-stage ovarian cancer.⁴² Human epididymis protein 4 is another serum tumor marker that can be useful for early epithelial ovarian cancer detection (sensitivity, 65%-83%; specificity, 78%-99%).⁴³ Elevated levels of CA19-9 and

Box 2. Diagnostic Workup for Epithelial Ovarian Cancer

- Cancer history (personal and family)
 - Hereditary disorders: hereditary breast and ovarian cancer syndrome (*BRCA* gene variants), Lynch syndrome
 - Breast, ovarian, endometrial, colon, or pancreatic cancer in first- or second-degree relatives
- Clinical examination (including pelvic examination)
- Serum tumor markers
 - CA125
 - CA19-9 and CEA (for suspected [based on imaging findings] or confirmed mucinous histology)
- Imaging
 - Pelvic ultrasound by expert examiner (to characterize adnexal masses, especially in early-stage disease)
 - Computed tomography of the chest, abdomen, and pelvis with oral and intravenous contrast (for clinical staging and surgical planning)
 - Additional tools in selected cases
 - Abdominal diffusion-weighted magnetic resonance imaging, laparoscopy, or mini laparotomy (to assess extent of peritoneal carcinomatosis and resectability)
 - Total-body 18F-fluorodeoxyglucose-positron emission tomography (to characterize suspicious distant lymph nodes and extra-abdominal sites)
 - Endoscopy (esophagogastroduodenoscopy and colonoscopy) if suspected (based on elevated CA19-9, elevated CEA, CA125/CEA ratio ≤ 25 , and/or imaging findings) or confirmed mucinous histology, to exclude primary gastrointestinal tumors
 - Pathological examination of adequate tumor sample from diagnostic core biopsy or surgical specimen^a
 - Cytological assessment of ascites or pleural effusion, if present
 - Genetic and molecular testing
 - Germline testing for *BRCA1/2* and other ovarian cancer susceptibility genes for all patients with epithelial ovarian cancer (to screen for hereditary breast and ovarian cancer syndrome and inform treatment decisions). In patients without a germline pathogenic *BRCA1/2* variant, somatic tumor testing for *BRCA1/2* should be performed to guide treatment decisions.
 - Tumor testing for homologous recombination deficiency for advanced high-grade serous or endometrioid ovarian cancers (to inform treatment decisions)
 - Tumor mismatch repair system testing—specifically, immunohistochemistry and microsatellite instability testing—for patients with endometrioid, clear cell, or mucinous subtypes (to screen for Lynch syndrome)

Abbreviations: CA125, carbohydrate antigen 125; CA19-9, carbohydrate antigen 19-9; CEA, carcinoembryonic antigen.

^a Preoperative biopsy should be avoided in presumed early-stage disease to prevent spilling malignant cells into the peritoneal cavity.

CEA, associated with mucinous gastrointestinal cancers, may be detected in patients with mucinous ovarian tumors. Primary ovarian mucinous cancers are rare, so upper and lower endoscopy should be performed to evaluate for primary gastrointestinal tumor metastatic to the ovary. A CA125/CEA ratio greater than 25 is suggestive of a primary mucinous ovarian tumor instead of gastrointestinal metastasis to the ovary.^{2,4,41}

Biopsy

Preoperative biopsy (eg, fine-needle aspiration) is contraindicated in patients with presumed early-stage ovarian cancer because it may

lead to tumor rupture and spillage of malignant cells into the peritoneal cavity. Intraoperative frozen section of a surgical specimen is usually performed to make the pathological diagnosis of ovarian cancer.^{4,44} For patients with advanced-stage disease not suitable for primary surgery, core biopsy (preferred) or fine-needle aspiration preoperative tissue analysis is required to confirm the diagnosis of ovarian cancer and guide chemotherapy selection. For patients with advanced-stage disease who are eligible for initial surgery without preoperative chemotherapy, diagnostic intraoperative biopsy suffices. For patients with ascites or pleural effusion, paracentesis or thoracentesis can provide cytological diagnosis and symptom relief.^{2,4}

Genetic and Molecular Testing


The NCCN and the American Society of Clinical Oncology guidelines recommend germline blood testing for all patients with epithelial ovarian cancer to screen for inherited *BRCA1/2* variants.^{10,13,45} Negative germline test findings should prompt surgical specimen evaluation for tumor-specific acquired (somatic) *BRCA1/2* variants. Germline and somatic testing inform treatment and counseling of patients and at-risk relatives (Box 2).^{10,13,45} Testing for HRD in advanced-stage, high-grade serous or endometrioid ovarian cancer also guides treatment,⁴⁶ with HRD positivity associated with a higher response to platinum-based chemotherapy and treatment with DNA repair inhibitors such as PARP inhibitors.⁴² Mismatch repair tumor testing is recommended by NCCN guidelines for endometrioid, clear cell, or mucinous subtypes to screen for Lynch syndrome.^{4,13}

Treatment of Early-Stage Ovarian Cancer

First-line therapy for presumed early-stage ovarian cancer is surgery to resect and stage the tumor, followed by adjuvant platinum-based chemotherapy (Figure 2). Surgery for most patients with early-stage ovarian cancer includes (1) en bloc resection of the involved ovary and fallopian tube to avoid spillage of cancer cells into the peritoneum during surgery; (2) hysterectomy and salpingo-oophorectomy of the contralateral ovary and fallopian tube; (3) retroperitoneal (pelvic and para-aortic) lymphadenectomy; (4) omentectomy; (5) random peritoneal biopsies; and (6) peritoneal fluid cytology. Appendectomy may also be performed for some patients with mucinous ovarian cancer.^{4,47} Intraoperative staging is important because approximately 30% of patients with presumed early-stage ovarian cancer by imaging have occult metastatic disease on definitive pathological examination of retroperitoneal lymph nodes, omentum, and peritoneal surface biopsies.^{4,47}

Patients with stage I disease who are considering having children in the future may undergo fertility-sparing surgery, consisting of unilateral salpingo-oophorectomy (preserving the uterus and contralateral ovary/fallopian tube without ovarian cancer).^{2,4} Cryopreservation of oocytes before bilateral salpingo-oophorectomy may allow future assisted reproductive approaches, such as in vitro fertilization and uterine embryo transfer. Consultation with an oncofertility specialist preoperatively is recommended. In a multi-institutional retrospective study of 211 women undergoing

Figure 2. Primary Treatment for Early-Stage and Advanced-Stage Ovarian Cancer



ASA indicates American Society of Anesthesiologists; AUC, area under the curve; CA125, carbohydrate antigen 125; ECOG-PS, Eastern Cooperative Oncology Group performance status; HIPEC, hyperthermic intraperitoneal chemotherapy; and PARP, poly-adenosine diphosphate ribose polymerase.

^aConsider neoadjuvant chemotherapy also if recent (<6 months) venous thromboembolism, myocardial infarction, stent, or laparotomy.

^bAn alternative weekly chemotherapy regimen of carboplatin AUC 2 and paclitaxel, 60 mg/m², is a reasonable option for frail patients who cannot tolerate the every-3-week schedule.

^cBevacizumab should be stopped 6 to 8 weeks before surgery and reinitiated 4 to 8 weeks after surgery due to wound-healing issues.

fertility-sparing surgery for unilateral epithelial ovarian cancer (stage IA, n = 126; stage IC, n = 85), 5-year overall and recurrence-free survival varied by subtype, with the highest rates (100% and 97.8%) for stage IA grade 1 to 2 disease and lower rates (66.7% and 66.0%) for stage IC grade 3 disease.⁴⁸

The benefit of adjuvant chemotherapy in early-stage ovarian cancer depends on stage, histotype, and tumor grade, and decisions about adjuvant chemotherapy should involve considerations

of age, comorbidities, performance status, and possible adverse events (Table; eTable 2 in the Supplement).^{49–54,65,67,68} A Cochrane meta-analysis of 5 RCTs including 1277 women with early-stage ovarian cancer found that adjuvant platinum-based chemotherapy improved 10-year overall survival compared with no chemotherapy (hazard ratio, 0.72; 95% CI, 0.57–0.92; absolute data not available).⁹⁷ Among patients with high-risk disease (stage IA grade 3, stage IB-IC grade 2–3, stage II, or any clear cell ovarian cancer), those who

Table. Summary of Evidence on Standard Treatment Strategies for Ovarian Cancer

Setting of therapy	Selected phase 3 randomized clinical trials	No. of randomized patients	Survival data ^a	Common serious adverse events ^{a,b}	Conclusions
Early-stage ovarian cancer					
Adjuvant systemic chemotherapy after surgery	ACTION, ^{49,50} ICON1, ^{51,52} GOG-157, ⁵³ ICON3 ⁵⁴	1765	10-Year overall survival: 73%-82% for adjuvant platinum-based chemotherapy vs 64%-76% for observation	Neutropenia (86%), thrombocytopenia (21%), anemia (7%), neurotoxicity (grade ≥2: 18%-29%), nausea/vomiting (9%-10%), allergy (3%), alopecia (grade 2: 64%-80%)	Adjuvant platinum-based chemotherapy (carboplatin and paclitaxel) is well tolerated and improves long-term overall survival
Advanced-stage ovarian cancer					
Cytoreductive surgery	EORTC 55971, ⁵⁵ CHORUS, ⁵⁶ JCOG0602, ⁵⁷ SCORPION ⁵⁸	1654	Median overall survival: 24-44 months for interval cytoreductive surgery vs 23-49 months for primary cytoreductive surgery	28-Day mortality (<1% vs 1%-6%), hemorrhage (4%-6% vs 3%-7%), venous thrombosis (0%-3% vs 2%-5%), infection (1%-3% vs 1%-8%), gastrointestinal fistula (<1% vs 1%-3%)	Survival with neoadjuvant chemotherapy followed by interval cytoreductive surgery is not inferior to primary cytoreductive surgery, which, however, carries a higher risk of perioperative complications
First-line systemic chemotherapy	ICON3, ⁵⁴ GOG-158, ⁵⁹ AGO-OVAR-3, ⁶⁰ SCOTROC1, ⁶¹ JGOG3016, ^{62,63} MITO-2, ⁶⁴ MITO-7, ⁶⁵ GOG-262, ⁶⁶ ICON8 ^{67,68}	8847	Median overall survival: 36-57 months for carboplatin-paclitaxel vs 35-49 months for other platinum-based regimens	Neutropenia (11%-88% vs 3%-94%), thrombocytopenia (2%-39% vs 1%-16%), anemia (1%-44% vs 1%-11%), neurotoxicity (3%-30% vs 1%-13%), nausea/vomiting (3%-10% vs 10%-23%), allergy (2%-3% vs 3%), alopecia (grade 2: 60%-80% vs 5%-76%)	Given the favorable toxicity profile and efficacy, carboplatin-paclitaxel is the standard first-line chemotherapy
Bevacizumab concurrent with chemotherapy and as maintenance therapy	ICON7, ^{69,70} GOG-0218 ^{71,72}	3401	Median PFS: 14-24 months for chemotherapy plus bevacizumab vs 10-22 months for chemotherapy alone	Hypertension (18%-23% vs 2%-7%), proteinuria (1%-2% vs <1%), gastrointestinal perforation (1% vs <1%), bleeding (1%-2% vs <1%), venous thromboembolism (7% vs 6%), wound dehiscence (1%-3% vs <1%-3%)	Addition of bevacizumab to platinum-based chemotherapy followed by bevacizumab maintenance therapy prolongs PFS by 2-4 months
PARP inhibitor (olaparib or niraparib) maintenance therapy after response to platinum-based chemotherapy	SOLO1, ^{73,74} PAOLA-1, ^{75,76} PRIMA ^{77,78}	1930	• BRCA variants: 7-year overall survival: 67% for olaparib vs 47% for placebo • HRD positive: median overall survival: 75 months for olaparib plus bevacizumab vs 57 months for bevacizumab • HRD negative: median PFS: 8.4 months for niraparib vs 5.4 months for placebo	Neutropenia (6%-15%), anemia (17%-31%), thrombocytopenia (1%-29%), fatigue (2%-5%), vomiting (1%), diarrhea (2%-3%), abdominal pain (1%-2%), hypertension (7%)	• Olaparib improves long-term overall survival in patients with BRCA variants • Olaparib plus bevacizumab improves overall survival in patients with HRD-positive ovarian cancer • Niraparib improves PFS regardless of BRCA or HRD status
Platinum-sensitive recurrent ovarian cancer					
Secondary cytoreductive surgery	GOG-213, ⁷⁹ DESKTOP III, ⁸⁰ SOC-1 ^{81,82}	1249	Median overall survival: 51-58 months for secondary cytoreductive surgery followed by platinum-based chemotherapy vs 46-65 months for platinum-based chemotherapy	Surgical morbidity at 30 days was 5%-9%; patient-reported quality of life did not differ between the 2 groups	Secondary cytoreductive surgery before platinum-based chemotherapy may be considered if complete resection of macroscopic disease is achievable
Second-line systemic chemotherapy	ICON4, ⁸³ Pfisterer et al, ⁸⁴ CALYPSO, ^{85,86} OCEANS, ^{87,88} GOG-213, ⁸⁹ AGO-OVAR-2, ^{21,90} bevacizumab	3974	Median overall survival: 18-42 months for platinum-based doublets (carboplatin plus paclitaxel, gemcitabine, or pegylated liposomal doxorubicin) with or without bevacizumab	Neutropenia (70%), anemia (27%), thrombocytopenia (35%), fatigue (2%), vomiting (3%), neurotoxicity (grade ≥2: 20%)	Platinum-based doublet chemotherapy with or without bevacizumab is the standard treatment for platinum-sensitive recurrent ovarian cancer
PARP inhibitor (olaparib, niraparib, or rucaparib) maintenance therapy	SOLO2, ^{91,92} NOVA, ⁹³ ARIEL-3 ⁹⁴	1412	BRCA variants: median overall survival: 41-52 months for PARP inhibitor vs 38-39 months for placebo	Neutropenia (5%-20%), anemia (19%-25%), thrombocytopenia (1%-34%), hypertension (8%), fatigue (4%-8%), vomiting (2%-4%), diarrhea (1%), abdominal pain (1%-3%)	Second-line PARP inhibitor maintenance therapy improves overall survival in patients with BRCA variants who did not previously receive a PARP inhibitor

(continued)

Table. Summary of Evidence on Standard Treatment Strategies for Ovarian Cancer (continued)

Setting of therapy	Selected phase 3 randomized clinical trials	No. of randomized patients	Survival data ^a	Common serious adverse events ^{a,b}	Conclusions
Platinum-resistant recurrent ovarian cancer					
Second-line systemic chemotherapy	AURELIA ⁹⁵	361	Median PFS: 6.7 months for single-agent chemotherapy (pegylated liposomal doxorubicin, weekly paclitaxel, or topotecan) plus bevacizumab vs 3.4 months for chemotherapy alone	Hypertension (grade ≥2: 20% vs 7%), proteinuria (2% vs 0%), gastrointestinal perforation (2% vs 0%), fistula/abscess (2% vs 0%), thromboembolism (5% vs 4%)	Non-platinum-based single-agent chemotherapy plus bevacizumab is the traditional standard treatment for platinum-resistant recurrent ovarian cancer
Mirvetuximab soravtansine (anti-folate receptor α antibody-drug conjugate)	MIRASOL ⁹⁶	453	Median overall survival: 16.5 months for mirvetuximab vs 12.7 months for single-agent chemotherapy	Blurred vision (7.8% vs 0%), keratopathy (9.2% vs 0%), dry eye (3.2% vs 0%), neutropenia (0.9% vs 1.4%), anemia (0.9% vs 1.1%), abdominal pain (2.8% vs 1.4%)	Mirvetuximab is the new standard treatment for patients with platinum-resistant recurrent ovarian cancer with high folate receptor α expression

^b Grade 3 or higher according to Common Terminology Criteria for Adverse Events version 5.0 unless otherwise specified.

Abbreviations: HRD, homologous recombination deficiency; PARP, poly-adenosine diphosphate ribose polymerase; PFS, progression-free survival.

^a Survival data rates and adverse event rates across randomized clinical trials are reported as ranges.

received adjuvant chemotherapy had improved 10-year overall survival compared with those not treated with adjuvant chemotherapy (hazard ratio, 0.52; 95% CI, 0.33-0.81; absolute data not available).⁹⁷


Treatment of Advanced-Stage Ovarian Cancer

Treatment of advanced-stage epithelial ovarian cancer typically involves a combination of cytoreductive surgery, systemic chemotherapy, and individualized maintenance therapy (Figure 2).^{2,4} For patients at low risk of perioperative complications and resectable advanced-stage ovarian cancer based on imaging, primary cytoreductive surgery (hysterectomy with bilateral salpingo-oophorectomy and omentectomy) is recommended before adjuvant platinum-based chemotherapy.⁹⁸⁻¹⁰⁰ Patients with ovarian cancer involving other organs may require resection of enlarged retroperitoneal pelvic and para-aortic lymph nodes, spleen, bowel, and diaphragm.⁹⁸⁻¹⁰¹

Patients with advanced-stage ovarian cancer at high risk of perioperative complications or unlikely to achieve complete cytoreduction at primary surgery should receive neoadjuvant chemotherapy followed by interval cytoreductive surgery and adjuvant systemic chemotherapy (Table; eTable 2 in the [Supplement](#)).^{55-58,102} A meta-analysis of 4 phase 3 RCTs including 1692 patients with advanced-stage ovarian cancer found that those who underwent interval cytoreductive surgery (n = 847) had similar overall survival (hazard ratio, 0.96; 95% CI, 0.86-1.08) and lower rates of postoperative complications (relative risk, 0.22; 95% CI, 0.13-0.38)¹⁰³ compared with those who underwent primary cytoreductive surgery (n = 845).¹⁰³ For patients with stage III ovarian cancer, hyperthermic intraperitoneal chemotherapy, which involves delivery of heated platinum-based chemotherapy into the abdominal cavity during surgery, may be offered during interval cytoreductive surgery (eTable 2 in the [Supplement](#)).^{102,104-107}

First-line systemic chemotherapy for advanced-stage ovarian cancer consists of carboplatin plus paclitaxel every 3 weeks for 6 cycles (Table; eTable 2 in the [Supplement](#)).^{4,59-68,108,109} Patients undergoing primary cytoreductive surgery receive 6 cycles of chemotherapy, starting within 6 weeks after surgery. Patients undergoing interval cytoreductive surgery typically receive 3 to 4 chemotherapy cycles prior to surgery and 2 to 3 after surgery.¹⁰² A weekly regimen of carboplatin and paclitaxel delivering the same cumulative dose as the every-3-week schedule is a reasonable alternative for patients who are unable to tolerate the standard schedule.^{65,67,68} Two large phase 3 RCTs (GOG-0218 and ICON7) demonstrated that adding bevacizumab to chemotherapy followed by bevacizumab maintenance therapy for up to 22 cycles (15 months) in patients with advanced-stage ovarian cancer prolonged mean progression-free survival (PFS) by 2 to 4 months vs chemotherapy alone (GOG-0218, 14.1 months vs 10.3 months; median follow-up, 17 months; ICON7, 24.1 months vs 22.4 months; median follow up, 28 months) (Table; eTable 2 in the [Supplement](#)).^{69,71} When these 2 trials were combined in a meta-analysis of 3401 patients, the bevacizumab group had improved PFS (hazard ratio, 0.72; 95% CI, 0.65-0.81; absolute data not available) compared with chemotherapy alone, although with no overall survival improvement.¹¹⁰ An exploratory analysis of a predefined subgroup of 502 high-risk patients (stage IV,

Figure 3. Treatment Algorithm for Recurrent Ovarian Cancer



inoperable stage III, or >1-cm residual disease after surgery for stage III ovarian cancer) in ICON7 showed improvement in mean overall survival in the bevacizumab group compared with chemotherapy alone (39.3 months vs 34.5 months; hazard ratio, 0.78; 95% CI, 0.63-0.97).⁷⁰

Bevacizumab and/or PARP inhibitor (olaparib or niraparib) maintenance therapy after primary treatment may be considered for patients with advanced-stage ovarian cancer depending on their response to chemotherapy, histology, *BRCA*/HRD status, and comorbidities (Table; eTable 2 in the Supplement).^{69-78,111-113} PARP inhibitors are guideline recommended for maintenance therapy in patients with high-grade serous or endometrioid ovarian cancer who respond to platinum-based chemotherapy, especially in *BRCA* variant or HRD-positive tumors.^{2,4} PARP inhibitors are administered orally, with olaparib dosed at 300 mg twice daily for 2 years and niraparib dosed at 200 mg to 300 mg once daily for 3 years.⁶ Olaparib provided a 7-year overall survival benefit compared with placebo (67.0% vs 46.5%; median overall survival, not reached vs 75.2 months; hazard ratio, 0.55; 95% CI, 0.40-0.76) in a phase 3 RCT of 391 patients with *BRCA* variant high-grade advanced-stage ovarian cancer (eTable 2 in the Supplement).^{73,74} In another phase 3 trial of 806 patients with advanced-stage ovarian cancer responding to platinum-based chemotherapy and bevacizumab, addition of olaparib to bevacizumab improved 5-year overall survival in patients with HRD-positive tumors (including *BRCA* variants) compared with placebo plus bevacizumab (65.5% vs 48.4%; median overall survival, 75.2 months vs 57.3 months; hazard ratio, 0.62; 95% CI, 0.45-0.85). Five-year overall survival rates were 73.2% vs 53.8% (median overall survival, 75.2 months vs 66.9 months; hazard ratio, 0.60; 95% CI, 0.39-0.93) for those with *BRCA* variants and 54.7% vs 44.2% (median overall survival, not reached vs 52.0 months; hazard ratio, 0.71; 95% CI, 0.45-1.13) for those without *BRCA* variants (eTable 2 in the Supplement).^{75,76} Three other trials investigating PARP inhibitor maintenance therapy found that PFS was

highest among patients with *BRCA* variant tumors followed by HRD-positive tumors and lowest for those with HRD-negative tumors (eTable 2 in the Supplement).^{77,111,112}

Prognosis and Surveillance

Overall 5-year survival rates are 70% to 95% for patients treated for early-stage ovarian cancer and 10% to 40% for those with advanced-stage disease.¹¹⁴ More than 80% of patients with advanced-stage ovarian cancer have initial complete remission (no disease on post-treatment imaging). PARP inhibitor maintenance therapy after platinum-based chemotherapy improves 5-year overall survival rates to approximately 70% in patients with *BRCA* variant or HRD-positive advanced-stage ovarian cancer.^{74,76}

There are currently no evidence-based treatment follow-up protocols, so surveillance should be individualized based on stage of disease, histology, genetic variants, and radiological and serum tumor marker response to primary treatment. Patients with complete remission after primary treatment should receive clinical reevaluation and CA125 testing (if initially elevated) every 2 to 4 months for the first 2 years, every 3 to 6 months during years 3 to 5, and then annually. Computed tomographic imaging should be individualized based on recurrence risk or clinical suspicion of recurrence (eg, abdominal pain, bloating, weight loss, increasing CA125 level).^{2,4}

Ovarian Cancer Relapse

Approximately 75% of patients with advanced-stage ovarian cancer and 10% to 30% of patients with early-stage ovarian cancer relapse within 2 years.^{2,4} Decisions about additional therapy, enrolling in clinical trials, or initiating palliative care should be made on an individual basis (Figure 3). Recurrent disease treatment depends on

the platinum-free interval—the time from the last dose of platinum-based chemotherapy to recurrence—which predicts response to subsequent platinum therapy. At first relapse, approximately 80% of patients have platinum-sensitive disease (platinum-free interval ≥ 6 months) and 20% have platinum-resistant disease (platinum-free interval < 6 months).¹¹⁵ With each remission and recurrence, platinum-free intervals shorten, and about 60% to 75% of patients with recurrent disease develop platinum resistance.¹¹⁵ Patients with recurrent ovarian cancer should undergo tumor testing for folate receptor α (FR α [FOLR1]) and *ERBB2* (previously *HER2/neu*) to identify potential targets for therapy.

Platinum-Sensitive Ovarian Cancer

For patients with platinum-sensitive recurrent ovarian cancer, platinum-based chemotherapy (carboplatin plus pegylated liposomal doxorubicin, gemcitabine, or paclitaxel) with or without bevacizumab is recommended (Table; eTable 2 in the Supplement).⁸³⁻⁹⁰ The median PFS with this treatment is 8 to 13 months.¹¹⁵ Secondary cytoreductive surgery prior to chemotherapy may be considered for platinum-sensitive disease if imaging and performance status indicate potentially resectable disease (Table; eTable 2 in the Supplement).⁷⁹⁻⁸²

Clinical trial data support use of maintenance therapy with a PARP inhibitor (olaparib, niraparib, or rucaparib) for recurrent disease in PARP inhibitor-naïve patients with *BRCA1/2* gene variants who responded to platinum-based chemotherapy (Table).^{91-94,116,117} A phase 3 RCT including 295 patients with *BRCA* gene variants and platinum-sensitive recurrent ovarian cancer showed that olaparib maintenance provided a median overall survival benefit of 12.9 months compared with placebo (51.7 months vs 38.8 months; hazard ratio, 0.74; 95% CI, 0.54-1.00).^{91,92} Niraparib and rucaparib have similar results (eTable 2 in the Supplement).^{93,94,116,117}

For patients with recurrent ovarian cancer without a *BRCA* variant or who received PARP inhibitor as first-line treatment, bevacizumab maintenance can be considered.¹¹⁸ A phase 3 RCT including 484 patients with first platinum-sensitive recurrence reported that adding bevacizumab to platinum-based chemotherapy followed by bevacizumab maintenance improved median PFS (12.4 months vs 8.4 months; hazard ratio, 0.48; 95% CI, 0.38-0.61) compared with chemotherapy alone (eTable 2 in the Supplement).^{87,88}

Platinum-Resistant Ovarian Cancer

Platinum-resistant recurrent ovarian cancer may be treated with single-agent non-platinum-based chemotherapy (pegylated liposomal doxorubicin, gemcitabine, weekly paclitaxel, or topotecan), associated with a median PFS of 3 to 9 months.^{95,115} Combining bevacizumab with chemotherapy improved PFS compared with chemotherapy alone in a phase 3 RCT of 361 patients with platinum-resistant recurrent ovarian cancer (6.7 months vs 3.4 months; hazard ratio, 0.48; 95% CI, 0.38-0.61) (Table; eTable 2 in the Supplement).⁹⁵

For the 30% to 35% of patients with platinum-resistant ovarian cancers exhibiting high FR α expression who have received 1 to 3 rounds of any type of anticancer therapy, the US Food and Drug Administration (FDA) has approved an antibody-drug conjugate targeting FR α , mirvetuximab soravtansine.^{96,119} In a phase 3 RCT of 453 patients, mirvetuximab provided a median overall survival benefit compared with conventional chemotherapy (16.5 months vs 12.7 months; hazard ratio, 0.67; 95% CI, 0.50-0.89) (Table; eTable 2 in the Supplement).⁹⁶ Another recently FDA-approved antibody-drug conjugate is trastuzumab deruxtecan, which targets *ERBB2* and provides a treatment option for patients with *ERBB2*-expressing ovarian cancer (eTable 2 in the Supplement).¹²⁰

Palliative Care

The NCCN guidelines recommend providing supportive care for all patients with ovarian cancer, including referral for palliative care assessment for those with advanced-stage or recurrent disease.⁴ Palliative care clinicians provide comprehensive multimodal symptom management, addressing pain, mental health, sexual health, and menopausal symptoms, and can facilitate transition to hospice care.¹²¹ Use of palliative care optimizes quality of life and may improve survival.^{121,122} Although not specific to ovarian cancer, outpatient palliative care for advanced-stage cancers has been shown to increase 1-year survival by 14% (56% vs 42%; $P < .001$) and median overall survival by 4.6 months (14.6 months vs 10.0 months) compared with usual care.¹²³

Limitations

This Review has several limitations. First, it is not a systematic review, and the quality of included studies was not formally evaluated. Second, this Review did not include less common epithelial and nonepithelial ovarian cancers. Third, treatments, outcomes, and prognostic estimates were derived from trials conducted primarily in North America and Europe and may not be generalizable to lower-income countries.

Conclusions

Approximately 21 000 women are diagnosed with ovarian cancer annually in the US, and approximately 80% have advanced-stage ovarian cancer at diagnosis. First-line treatment of early-stage ovarian cancer is surgery and adjuvant platinum-based chemotherapy. Treatment of advanced-stage ovarian cancer includes cytoreductive surgery, platinum-based chemotherapy, and targeted maintenance therapies such as bevacizumab and/or PARP inhibitors.

ARTICLE INFORMATION

Accepted for Publication: May 21, 2025.

Published Online: July 21, 2025.
doi:10.1001/jama.2025.9495

Conflict of Interest Disclosures: Dr Cliby reported receipt of a National Institutes of Health SPOR

grant in ovarian cancer. No other disclosures were reported.

Submissions: We encourage authors to submit papers for consideration as a Review. Please contact Kristin Walter, MD, at kristin.walter@jamanetwork.org.

REFERENCES

1. Siegel RL, Kratzer TB, Giaquinto AN, Sung H, Jemal A. Cancer statistics, 2025. *CA Cancer J Clin*. 2025;75(1):10-45. doi:10.3322/caac.21871
2. González-Martín A, Harter P, Leary A, et al; ESMO Guidelines Committee. Newly diagnosed and relapsed epithelial ovarian cancer: ESMO clinical

- practice guideline for diagnosis, treatment and follow-up. *Ann Oncol*. 2023;34(10):833-848. doi:10.1016/j.annonc.2023.07.011
3. Huepenbecker SP, Sun CC, Fu S, et al. Factors impacting the time to ovarian cancer diagnosis based on classic symptom presentation in the United States. *Cancer*. 2021;127(22):4151-4160. doi:10.1002/cncr.33829
 4. National Comprehensive Cancer Network. Ovarian cancer (version 1.2025). Accessed April 5, 2025. https://www.nccn.org/professionals/physician_gls/pdf/ovarian.pdf
 5. Female genital tumours. In: WHO Classification of Tumours Editorial Board, ed. *World Health Organization Classification of Tumours*. 5th ed. IARC Press; 2020.
 6. Caruso G, Tomao F, Parma G, et al. Poly (ADP-ribose) polymerase inhibitors (PARPi) in ovarian cancer: lessons learned and future directions. *Int J Gynecol Cancer*. 2023;33(4):431-443. doi:10.1136/ijgc-2022-004149
 7. Bray F, Laversanne M, Sung H, et al. Global cancer statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*. 2024;74(3):229-263. doi:10.3322/caac.21834
 8. Cabasag CJ, Fagan PJ, Ferlay J, et al. Ovarian cancer today and tomorrow: a global assessment by world region and Human Development Index using GLOBOCAN 2020. *Int J Cancer*. 2022;151(9):1535-1541. doi:10.1002/ijc.34002
 9. Kotsopoulos J, Gronwald J, Karlan B, et al; Hereditary Ovarian Cancer Clinical Study Group. Age-specific ovarian cancer risks among women with a *BRCA1* or *BRCA2* mutation. *Gynecol Oncol*. 2018;150(1):85-91. doi:10.1016/j.jgyno.2018.05.011
 10. Daly MB, Pal T, Maxwell KN, et al. NCCN guidelines insights: genetic/familial high-risk assessment: breast, ovarian, and pancreatic, version 2.2024. *J Natl Compr Canc Netw*. 2023;21(10):1000-1010. doi:10.6004/jnccn.2023.0051
 11. Walsh T, Casadei S, Lee MK, et al. Mutations in 12 genes for inherited ovarian, fallopian tube, and peritoneal carcinoma identified by massively parallel sequencing. *Proc Natl Acad Sci U S A*. 2011;108(44):18032-18037. doi:10.1073/pnas.1115052108
 12. Bonadona V, Bonaïti B, Olschwang S, et al; French Cancer Genetics Network. Cancer risks associated with germline mutations in *MLH1*, *MSH2*, and *MSH6* genes in Lynch syndrome. *JAMA*. 2011;305(22):2304-2310. doi:10.1001/jama.2011.743
 13. Konstantinopoulos PA, Norquist B, Lacchetti C, et al. Germline and somatic tumor testing in epithelial ovarian cancer: ASCO guideline. *J Clin Oncol*. 2020;38(11):1222-1245. doi:10.1200/JCO.19.02960
 14. Barnard ME, Farland LV, Yan B, et al. Endometriosis typology and ovarian cancer risk. *JAMA*. 2024;332(6):482-489. doi:10.1001/jama.2024.9210
 15. Tanha K, Mottaghi A, Nojomi M, et al. Investigation on factors associated with ovarian cancer: an umbrella review of systematic review and meta-analyses. *J Ovarian Res*. 2021;14(1):153. doi:10.1186/s13048-021-00911-z
 16. Karnezis AN, Cho KR, Gilks CB, Pearce CL, Huntsman DG. The disparate origins of ovarian cancers: pathogenesis and prevention strategies. *Nat Rev Cancer*. 2017;17(1):65-74. doi:10.1038/nrc.2016.113
 17. Konstantinopoulos PA, Ceccaldi R, Shapiro GI, D'Andrea AD. Homologous recombination deficiency: exploiting the fundamental vulnerability of ovarian cancer. *Cancer Discov*. 2015;5(11):1137-1154. doi:10.1158/2159-8290.CD-15-0714
 18. Walker JL, Powell CB, Chen LM, et al. Society of Gynecologic Oncology recommendations for the prevention of ovarian cancer. *Cancer*. 2015;121(13):2108-2120. doi:10.1002/cncr.29321
 19. Marchetti C, De Felice F, Palaia I, et al. Risk-reducing salpingo-oophorectomy: a meta-analysis on impact on ovarian cancer risk and all cause mortality in *BRCA1* and *BRCA2* mutation carriers. *BMC Womens Health*. 2014;14:150. doi:10.1186/s12905-014-0150-5
 20. van Bommel MH, Int'Hout J, Veldmate G, et al. Contraceptives and cancer risks in *BRCA1/2* pathogenic variant carriers: a systematic review and meta-analysis. *Hum Reprod Update*. 2023;29(2):197-217. doi:10.1093/humupd/dmac038
 21. Iodice S, Barile M, Rotmensz N, et al. Oral contraceptive use and breast or ovarian cancer risk in *BRCA1/2* carriers: a meta-analysis. *Eur J Cancer*. 2010;46(12):2275-2284. doi:10.1016/j.ejca.2010.04.018
 22. American College of Obstetricians and Gynecologists. ACOG committee opinion No. 774: opportunistic salpingectomy as a strategy for epithelial ovarian cancer prevention. *Obstet Gynecol*. 2019;133(4):e279-e284. doi:10.1097/AOG.0000000000003164
 23. Mor-Hadar D, Wilailak S, Berek J, McNally OM; FIGO Committee on Women's Cancer. FIGO position statement on opportunistic salpingectomy as an ovarian cancer prevention strategy. *Int J Gynaecol Obstet*. 2024;167(3):976-980. doi:10.1002/ijgo.15884
 24. Havrilesky LJ, Moorman PG, Lowery WJ, et al. Oral contraceptive pills as primary prevention for ovarian cancer: a systematic review and meta-analysis. *Obstet Gynecol*. 2013;122(1):139-147. doi:10.1097/AOG.0b013e318291c235
 25. Grossman DC, Curry SJ, Owens DK, et al; US Preventive Services Task Force. Screening for ovarian cancer: US Preventive Services Task Force recommendation statement. *JAMA*. 2018;319(6):588-594. doi:10.1001/jama.2017.21926
 26. Skates SJ, Greene MH, Buys SS, et al. Early detection of ovarian cancer using the risk of ovarian cancer algorithm with frequent CA125 testing in women at increased familial risk—combined results from two screening trials. *Clin Cancer Res*. 2017;23(14):3628-3637. doi:10.1158/1078-0432.CCR-15-2750
 27. Rosenthal AN, Fraser LS, Philpott S, et al; United Kingdom Familial Ovarian Cancer Screening Study Collaborators. Evidence of stage shift in women diagnosed with ovarian cancer during phase II of the United Kingdom Familial Ovarian Cancer Screening Study. *J Clin Oncol*. 2017;35(13):1411-1420. doi:10.1200/JCO.2016.69.9330
 28. Lai T, Kessel B, Ahn HJ, Terada KY. Ovarian cancer screening in menopausal females with a family history of breast or ovarian cancer. *J Gynecol Oncol*. 2016;27(4):e41. doi:10.3802/jgo.2016.27.e41
 29. Hermesen BB, Olivier RI, Verheijen RH, et al. No efficacy of annual gynaecological screening in *BRCA1/2* mutation carriers: an observational follow-up study. *Br J Cancer*. 2007;96(9):1335-1342. doi:10.1038/sj.bjc.6603725
 30. Goff BA, Mandel L, Muntz HG, Melancon CH. Ovarian carcinoma diagnosis. *Cancer*. 2000;89(10):2068-2075. doi:10.1002/1097-0142(200011)89:10<2068::AID-CNCR6>3.0.CO;2-Z
 31. Goff BA, Mandel LS, Melancon CH, Muntz HG. Frequency of symptoms of ovarian cancer in women presenting to primary care clinics. *JAMA*. 2004;291(22):2705-2712. doi:10.1001/jama.291.22.2705
 32. Peter E, Honnorat J, Desestret V. Paraneoplastic neurologic syndrome associated with gynecologic and breast malignancies. *Handb Clin Neurol*. 2024;200:409-417. doi:10.1016/B978-0-12-823912-4.00014-1
 33. Timmerman D, Planchamp F, Bourne T, et al. ESGO/ISUOG/IOTA/ESGE consensus statement on pre-operative diagnosis of ovarian tumors. *Int J Gynecol Cancer*. 2021;31(7):961-982. doi:10.1136/ijgc-2021-002565
 34. Fischerova D, Pinto P, Burgetova A, et al. Preoperative staging of ovarian cancer: comparison between ultrasound, CT and whole-body diffusion-weighted MRI (ISAAC study). *Ultrasound Obstet Gynecol*. 2022;59(2):248-262. doi:10.1002/uog.23654
 35. Tempany CM, Zou KH, Silverman SG, Brown DL, Kurtz AB, McNeil BJ. Staging of advanced ovarian cancer: comparison of imaging modalities—report from the Radiological Diagnostic Oncology Group. *Radiology*. 2000;215(3):761-767. doi:10.1148/radiology.215.3.r00jn25761
 36. Berek JS, Renz M, Kehoe S, Kumar L, Friedlander M. Cancer of the ovary, fallopian tube, and peritoneum: 2021 update. *Int J Gynaecol Obstet*. 2021;155(suppl 1):61-85. doi:10.1002/ijgo.13878
 37. Michielsen K, Dresen R, Vanslambrouck R, et al. Diagnostic value of whole body diffusion-weighted MRI compared to computed tomography for pre-operative assessment of patients suspected for ovarian cancer. *Eur J Cancer*. 2017;83:88-98. doi:10.1016/j.ejca.2017.06.010
 38. Fleming ND, Nick AM, Coleman RL, et al. Laparoscopic surgical algorithm to triage the timing of tumor reductive surgery in advanced ovarian cancer. *Obstet Gynecol*. 2018;132(3):545-554. doi:10.1097/AOG.0000000000002796
 39. Kang SK, Reinhold C, Atri M, et al; Expert Panel on Women's Imaging. ACR appropriateness criteria staging and follow-up of ovarian cancer. *J Am Coll Radiol*. 2018;15(5S):S198-S207. doi:10.1016/j.jacr.2018.03.015
 40. Yuan Y, Gu ZX, Tao XF, Liu SY. Computer tomography, magnetic resonance imaging, and positron emission tomography or positron emission tomography/computer tomography for detection of metastatic lymph nodes in patients with ovarian cancer: a meta-analysis. *Eur J Radiol*. 2012;81(5):1002-1006. doi:10.1016/j.ejrad.2011.01.112
 41. Wright AA, Bohlke K, Armstrong DK, et al. Neoadjuvant chemotherapy for newly diagnosed, advanced ovarian cancer: Society of Gynecologic Oncology and American Society of Clinical Oncology clinical practice guideline. *J Clin Oncol*. 2016;34(28):3460-3473. doi:10.1200/JCO.2016.68.6907
 42. Medeiros LR, Rosa DD, da Rosa MI, Bozzetti MC. Accuracy of CA 125 in the diagnosis of ovarian tumors: a quantitative systematic review. *Eur J*

Obstet Gynecol Reprod Biol. 2009;142(2):99-105. doi:10.1016/j.ejogrb.2008.08.011

43. Ferraro S, Braga F, Lanzoni M, Boracchi P, Biganzoli EM, Panteghini M. Serum human epididymis protein 4 vs carbohydrate antigen 125 for ovarian cancer diagnosis: a systematic review. *J Clin Pathol.* 2013;66(4):273-281. doi:10.1136/jclinpath-2012-201031
44. Vergote I, De Brabanter J, Fyles A, et al. Prognostic importance of degree of differentiation and cyst rupture in stage I invasive epithelial ovarian carcinoma. *Lancet.* 2001;357(9251):176-182. doi:10.1016/S0140-6736(00)03590-X
45. Tung N, Ricker C, Messersmith H, et al. Selection of germline genetic testing panels in patients with cancer: ASCO guideline. *J Clin Oncol.* 2024;42(21):2599-2615. doi:10.1200/JCO.24.00662
46. Miller RE, Leary A, Scott CL, et al. ESMO recommendations on predictive biomarker testing for homologous recombination deficiency and PARP inhibitor benefit in ovarian cancer. *Ann Oncol.* 2020;31(12):1606-1622. doi:10.1016/j.annonc.2020.08.2102
47. Colombo N, Sessa C, du Bois A, et al; ESMO-ESGO Ovarian Cancer Consensus Conference Working Group. ESMO-ESGO consensus conference recommendations on ovarian cancer: pathology and molecular biology, early and advanced stages, borderline tumours and recurrent disease. *Ann Oncol.* 2019;30(5):672-705. doi:10.1093/annonc/mdz062
48. Satoh T, Hatae M, Watanabe Y, et al. Outcomes of fertility-sparing surgery for stage I epithelial ovarian cancer: a proposal for patient selection. *J Clin Oncol.* 2010;28(10):1727-1732. doi:10.1200/JCO.2009.24.8617
49. Trimbos JB, Vergote I, Bolis G, et al; EORTC-ACTION Collaborators. Impact of adjuvant chemotherapy and surgical staging in early-stage ovarian carcinoma: European Organisation for Research and Treatment of Cancer-Adjuvant Chemotherapy in Ovarian Neoplasm trial. *J Natl Cancer Inst.* 2003;95(2):113-125. doi:10.1093/jnci/95.2.113
50. Trimbos B, Timmers P, Pecorelli S, et al. Surgical staging and treatment of early ovarian cancer: long-term analysis from a randomized trial. *J Natl Cancer Inst.* 2010;102(13):982-987. doi:10.1093/jnci/djq149
51. Colombo N, Guthrie D, Chiari S, et al; International Collaborative Ovarian Neoplasm (ICON) Collaborators. International Collaborative Ovarian Neoplasm trial 1: a randomized trial of adjuvant chemotherapy in women with early-stage ovarian cancer. *J Natl Cancer Inst.* 2003;95(2):125-132. doi:10.1093/jnci/95.2.125
52. Collinson F, Qian W, Fossati R, et al; ICON1 Collaborators. Optimal treatment of early-stage ovarian cancer. *Ann Oncol.* 2014;25(6):1165-1171. doi:10.1093/annonc/mdu116
53. Bell J, Brady MF, Young RC, et al; Gynecologic Oncology Group. Randomized phase III trial of three versus six cycles of adjuvant carboplatin and paclitaxel in early stage epithelial ovarian carcinoma: a Gynecologic Oncology Group study. *Gynecol Oncol.* 2006;102(3):432-439. doi:10.1016/j.ygyno.2006.06.013
54. International Collaborative Ovarian Neoplasm Group. Paclitaxel plus carboplatin versus standard

- chemotherapy with either single-agent carboplatin or cyclophosphamide, doxorubicin, and cisplatin in women with ovarian cancer: the ICON3 randomised trial. *Lancet.* 2002;360(9332):505-515. doi:10.1016/S0140-6736(02)09738-6
55. Vergote I, Coens C, Nankivell M, et al; EORTC; MRC CHORUS Study Investigators. Neoadjuvant chemotherapy versus debulking surgery in advanced tubo-ovarian cancers: pooled analysis of individual patient data from the EORTC 55971 and CHORUS trials. *Lancet Oncol.* 2018;19(12):1680-1687. doi:10.1016/S1470-2045(18)30566-7
 56. Kehoe S, Hook J, Nankivell M, et al. Primary chemotherapy versus primary surgery for newly diagnosed advanced ovarian cancer (CHORUS): an open-label, randomised, controlled, non-inferiority trial. *Lancet.* 2015;386(9990):249-257. doi:10.1016/S0140-6736(14)62223-6
 57. Onda T, Satoh T, Saito T, et al; Japan Clinical Oncology Group. Comparison of treatment invasiveness between upfront debulking surgery versus interval debulking surgery following neoadjuvant chemotherapy for stage III/IV ovarian, tubal, and peritoneal cancers in a phase III randomised trial: Japan Clinical Oncology Group Study JCOG0602. *Eur J Cancer.* 2016;64:22-31. doi:10.1016/j.ejca.2016.05.017
 58. Fagotti A, Ferrandina MG, Vizzelli G, et al. Randomized trial of primary debulking surgery versus neoadjuvant chemotherapy for advanced epithelial ovarian cancer (SCORPION-NCT01461850). *Int J Gynecol Cancer.* 2020;30(11):1657-1664. doi:10.1136/ijgc-2020-001640
 59. Ozols RF, Bundy BN, Greer BE, et al; Gynecologic Oncology Group. Phase III trial of carboplatin and paclitaxel compared with cisplatin and paclitaxel in patients with optimally resected stage III ovarian cancer: a Gynecologic Oncology Group study. *J Clin Oncol.* 2003;21(17):3194-3200. doi:10.1200/JCO.2003.02.153
 60. du Bois A, Lück HJ, Meier W, et al; Arbeitsgemeinschaft Gynäkologische Onkologie Ovarian Cancer Study Group. A randomized clinical trial of cisplatin/paclitaxel versus carboplatin/paclitaxel as first-line treatment of ovarian cancer. *J Natl Cancer Inst.* 2003;95(17):1320-1329. doi:10.1093/jnci/djg036
 61. Vasey PA, Jayson GC, Gordon A, et al; Scottish Gynaecological Cancer Trials Group. Phase III randomized trial of docetaxel-carboplatin versus paclitaxel-carboplatin as first-line chemotherapy for ovarian carcinoma. *J Natl Cancer Inst.* 2004;96(22):1682-1691. doi:10.1093/jnci/djh323
 62. Katsumata N, Yasuda M, Takahashi F, et al; Japanese Gynecologic Oncology Group. Dose-dense paclitaxel once a week in combination with carboplatin every 3 weeks for advanced ovarian cancer: a phase 3, open-label, randomised controlled trial. *Lancet.* 2009;374(9698):1331-1338. doi:10.1016/S0140-6736(09)61157-0
 63. Katsumata N, Yasuda M, Isonishi S, et al; Japanese Gynecologic Oncology Group. Long-term results of dose-dense paclitaxel and carboplatin versus conventional paclitaxel and carboplatin for treatment of advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer (JGOG 3016): a randomised, controlled, open-label trial. *Lancet Oncol.* 2013;14(10):1020-1026. doi:10.1016/S1470-2045(13)70363-2

64. Pignata S, Scambia G, Ferrandina G, et al. Carboplatin plus paclitaxel versus carboplatin plus pegylated liposomal doxorubicin as first-line treatment for patients with ovarian cancer: the MITO-2 randomized phase III trial. *J Clin Oncol.* 2011;29(27):3628-3635. doi:10.1200/JCO.2010.33.8566
65. Pignata S, Scambia G, Katsaros D, et al; Multicentre Italian Trials in Ovarian cancer (MITO-7); Groupe d'Investigateurs Nationaux pour l'Etude des Cancers Ovariens et du sein (GINECO); Mario Negri Gynecologic Oncology (MaNGO); European Network of Gynaecological Oncological Trial Groups (ENGOT-OV-10); Gynecologic Cancer InterGroup (GCIG) Investigators. Carboplatin plus paclitaxel once a week versus every 3 weeks in patients with advanced ovarian cancer (MITO-7): a randomised, multicentre, open-label, phase 3 trial. *Lancet Oncol.* 2014;15(4):396-405. doi:10.1016/S1470-2045(14)70049-X
66. Chan JK, Brady MF, Penson RT, et al. Weekly vs every-3-week paclitaxel and carboplatin for ovarian cancer. *N Engl J Med.* 2016;374(8):738-748. doi:10.1056/NEJMoa1505067
67. Clamp AR, James EC, McNeish IA, et al. Weekly dose-dense chemotherapy in first-line epithelial ovarian, fallopian tube, or primary peritoneal carcinoma treatment (ICON8): primary progression free survival analysis results from a GCIG phase 3 randomised controlled trial. *Lancet.* 2019;394(10214):2084-2095. doi:10.1016/S0140-6736(19)32259-7
68. Clamp AR, James EC, McNeish IA, et al. Weekly dose-dense chemotherapy in first-line epithelial ovarian, fallopian tube, or primary peritoneal cancer treatment (ICON8): overall survival results from an open-label, randomised, controlled, phase 3 trial. *Lancet Oncol.* 2022;23(7):919-930. doi:10.1016/S1470-2045(22)00283-2
69. Perren TJ, Swart AM, Pfisterer J, et al; ICON7 Investigators. A phase 3 trial of bevacizumab in ovarian cancer. *N Engl J Med.* 2011;365(26):2484-2496. doi:10.1056/NEJMoa1103799
70. Oza AM, Cook AD, Pfisterer J, et al; ICON7 Trial Investigators. Standard chemotherapy with or without bevacizumab for women with newly diagnosed ovarian cancer (ICON7): overall survival results of a phase 3 randomised trial. *Lancet Oncol.* 2015;16(8):928-936. doi:10.1016/S1470-2045(15)00086-8
71. Burger RA, Brady MF, Bookman MA, et al; Gynecologic Oncology Group. Incorporation of bevacizumab in the primary treatment of ovarian cancer. *N Engl J Med.* 2011;365(26):2473-2483. doi:10.1056/NEJMoa1104390
72. Tewari KS, Burger RA, Enserro D, et al. Final overall survival of a randomized trial of bevacizumab for primary treatment of ovarian cancer. *J Clin Oncol.* 2019;37(26):2317-2328. doi:10.1200/JCO.19.01009
73. Moore K, Colombo N, Scambia G, et al. Maintenance olaparib in patients with newly diagnosed advanced ovarian cancer. *N Engl J Med.* 2018;379(26):2495-2505. doi:10.1056/NEJMoa1810858
74. DiSilvestro P, Banerjee S, Colombo N, et al; SOLO1 Investigators. Overall survival with maintenance olaparib at a 7-year follow-up in patients with newly diagnosed advanced ovarian cancer and a BRCA mutation: the SOLO1/GOG 3004

- trial. *J Clin Oncol*. 2023;41(3):609-617. doi:10.1200/JCO.22.01549
75. Ray-Coquard I, Pautier P, Pignata S, et al; PAOLA-1 Investigators. Olaparib plus bevacizumab as first-line maintenance in ovarian cancer. *N Engl J Med*. 2019;381(25):2416-2428. doi:10.1056/NEJMoa1911361
76. Ray-Coquard I, Leary A, Pignata S, et al; PAOLA-1/ENGOT-OV25 Investigators. Olaparib plus bevacizumab first-line maintenance in ovarian cancer: final overall survival results from the PAOLA-1/ENGOT-OV25 trial. *Ann Oncol*. 2023;34(8):681-692. doi:10.1016/j.annonc.2023.05.005
77. González-Martín A, Pothuri B, Vergote I, et al; PRIMA/ENGOT-OV26/GOG-3012 Investigators. Niraparib in patients with newly diagnosed advanced ovarian cancer. *N Engl J Med*. 2019;381(25):2391-2402. doi:10.1056/NEJMoa1910962
78. Monk BJ, Barretina-Ginesta MP, Pothuri B, et al. Niraparib first-line maintenance therapy in patients with newly diagnosed advanced ovarian cancer: final overall survival results from the PRIMA/ENGOT-OV26/GOG-3012 trial. *Ann Oncol*. 2024;35(11):981-992. doi:10.1016/j.annonc.2024.08.2241
79. Coleman RL, Spirtos NM, Enserro D, et al. Secondary surgical cytoreduction for recurrent ovarian cancer. *N Engl J Med*. 2019;381(20):1929-1939. doi:10.1056/NEJMoa1902626
80. Harter P, Sehouli J, Vergote I, et al; DESKTOP III Investigators. Randomized trial of cytoreductive surgery for relapsed ovarian cancer. *N Engl J Med*. 2021;385(23):2123-2131. doi:10.1056/NEJMoa2103294
81. Shi T, Zhu J, Feng Y, et al. Secondary cytoreduction followed by chemotherapy versus chemotherapy alone in platinum-sensitive relapsed ovarian cancer (SOC-1): a multicentre, open-label, randomised, phase 3 trial. *Lancet Oncol*. 2021;22(4):439-449. doi:10.1016/S1470-2045(21)00006-1
82. Jiang R, Feng Y, Chen Y, et al; SOC-1 Investigators. Surgery versus no surgery in platinum-sensitive relapsed ovarian cancer: final overall survival analysis of the SOC-1 randomized phase 3 trial. *Nat Med*. 2024;30(8):2181-2188. doi:10.1038/s41591-024-02981-0
83. Parmar MKB, Ledermann JA, Colombo N, et al; ICON and AGO Collaborators. Paclitaxel plus platinum-based chemotherapy versus conventional platinum-based chemotherapy in women with relapsed ovarian cancer: the ICON4/AGO-OVAR-2.2 trial. *Lancet*. 2003;361(9375):2099-2106. doi:10.1016/S0140-6736(03)13718-X
84. Pfisterer J, Plante M, Vergote I, et al. Gemcitabine plus carboplatin compared with carboplatin in patients with platinum-sensitive recurrent ovarian cancer: an intergroup trial of the AGO-OVAR, the NCIC CTG, and the EORTC GCG. *J Clin Oncol*. 2006;24(29):4699-4707. doi:10.1200/JCO.2006.06.0913
85. Pujade-Lauraine E, Wagner U, Aavall-Lundqvist E, et al. Pegylated liposomal doxorubicin and carboplatin compared with paclitaxel and carboplatin for patients with platinum-sensitive ovarian cancer in late relapse. *J Clin Oncol*. 2010;28(20):3323-3329. doi:10.1200/JCO.2009.25.7519
86. Wagner U, Marth C, Largillier R, et al. Final overall survival results of phase III GCIG CALYPSO trial of pegylated liposomal doxorubicin and carboplatin vs paclitaxel and carboplatin in platinum-sensitive ovarian cancer patients. *Br J Cancer*. 2012;107(4):588-591. doi:10.1038/bjc.2012.307
87. Aghajanian C, Blank SV, Goff BA, et al. OCEANS: a randomized, double-blind, placebo-controlled phase III trial of chemotherapy with or without bevacizumab in patients with platinum-sensitive recurrent epithelial ovarian, primary peritoneal, or fallopian tube cancer. *J Clin Oncol*. 2012;30(17):2039-2045. doi:10.1200/JCO.2012.42.0505
88. Aghajanian C, Goff B, Nycum LR, Wang YV, Husain A, Blank SV. Final overall survival and safety analysis of OCEANS, a phase 3 trial of chemotherapy with or without bevacizumab in patients with platinum-sensitive recurrent ovarian cancer. *Gynecol Oncol*. 2015;139(1):10-16. doi:10.1016/j.ygyno.2015.08.004
89. Coleman RL, Brady MF, Herzog TJ, et al. Bevacizumab and paclitaxel-carboplatin chemotherapy and secondary cytoreduction in recurrent, platinum-sensitive ovarian cancer (NRG Oncology/Gynecologic Oncology Group study GOG-0213): a multicentre, open-label, randomised, phase 3 trial. *Lancet Oncol*. 2017;18(6):779-791. doi:10.1016/S1470-2045(17)30279-6
90. Pfisterer J, Shannon CM, Baumann K, et al; AGO-OVAR 2.21/ENGOT-OV 18 Investigators. Bevacizumab and platinum-based combinations for recurrent ovarian cancer: a randomised, open-label, phase 3 trial. *Lancet Oncol*. 2020;21(5):699-709. doi:10.1016/S1470-2045(20)30142-X
91. Pujade-Lauraine E, Ledermann JA, Selle F, et al; SOLO2/ENGOT-OV21 Investigators. Olaparib tablets as maintenance therapy in patients with platinum-sensitive, relapsed ovarian cancer and a *BRCA1/2* mutation (SOLO2/ENGOT-OV21): a double-blind, randomised, placebo-controlled, phase 3 trial. *Lancet Oncol*. 2017;18(9):1274-1284. doi:10.1016/S1470-2045(17)30469-2
92. Poveda A, Floquet A, Ledermann JA, et al; SOLO2/ENGOT-OV21 Investigators. Olaparib tablets as maintenance therapy in patients with platinum-sensitive relapsed ovarian cancer and a *BRCA1/2* mutation (SOLO2/ENGOT-OV21): a final analysis of a double-blind, randomised, placebo-controlled, phase 3 trial. *Lancet Oncol*. 2021;22(5):620-631. doi:10.1016/S1470-2045(21)00073-5
93. Mirza MR, Monk BJ, Herrstedt J, et al; ENGOT-OV16/NOVA Investigators. Niraparib maintenance therapy in platinum-sensitive, recurrent ovarian cancer. *N Engl J Med*. 2016;375(22):2154-2164. doi:10.1056/NEJMoa1611310
94. Coleman RL, Oza AM, Lorusso D, et al; ARIEL3 Investigators. Rucaparib maintenance treatment for recurrent ovarian carcinoma after response to platinum therapy (ARIEL3): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet*. 2017;390(10106):1949-1961. doi:10.1016/S0140-6736(17)32440-6
95. Pujade-Lauraine E, Hilpert F, Weber B, et al. Bevacizumab combined with chemotherapy for platinum-resistant recurrent ovarian cancer: the AURELIA open-label randomized phase III trial. *J Clin Oncol*. 2014;32(13):1302-1308. doi:10.1200/JCO.2013.51.4489
96. Moore KN, Angelergues A, Konecny GE, et al; Gynecologic Oncology Group Partners; European Network of Gynaecological Oncological Trial Groups. Mirvetuximab soravtansine in FRα-positive, platinum-resistant ovarian cancer. *N Engl J Med*. 2023;389(23):2162-2174. doi:10.1056/NEJMoa2309169
97. Lawrie TA, Winter-Roach BA, Heus P, Kitchener HC. Adjuvant (post-surgery) chemotherapy for early stage epithelial ovarian cancer. *Cochrane Database Syst Rev*. 2015;2015(12):CD004706. doi:10.1002/14651858.CD004706.pub5
98. du Bois A, Reuss A, Pujade-Lauraine E, Harter P, Ray-Coquard I, Pfisterer J. Role of surgical outcome as prognostic factor in advanced epithelial ovarian cancer: a combined exploratory analysis of 3 prospectively randomized phase 3 multicenter trials: by the Arbeitsgemeinschaft Gynaekologische Onkologie Studiengruppe Ovarialkarzinom (AGO-OVAR) and the Groupe d'Investigateurs Nationaux Pour les Etudes des Cancers de l'Ovaire (GINECO). *Cancer*. 2009;115(6):1234-1244. doi:10.1002/cncr.24149
99. Chi DS, Eisenhauer EL, Lang J, et al. What is the optimal goal of primary cytoreductive surgery for bulky stage IIIC epithelial ovarian carcinoma (EOC)? *Gynecol Oncol*. 2006;103(2):559-564. doi:10.1016/j.ygyno.2006.03.051
100. Bristow RE, Tomacruz RS, Armstrong DK, Trimble EL, Montz FJ. Survival effect of maximal cytoreductive surgery for advanced ovarian carcinoma during the platinum era: a meta-analysis. *J Clin Oncol*. 2002;20(5):1248-1259. doi:10.1200/JCO.2002.20.5.1248
101. Harter P, Sehouli J, Lorusso D, et al. A randomized trial of lymphadenectomy in patients with advanced ovarian neoplasms. *N Engl J Med*. 2019;380(9):822-832. doi:10.1056/NEJMoa1808424
102. Gaillard S, Lacchetti C, Armstrong DK, et al. Neoadjuvant chemotherapy for newly diagnosed, advanced ovarian cancer: ASCO guideline update. *J Clin Oncol*. 2025;43(7):868-891. doi:10.1200/JCO.24.02589
103. Coleridge SL, Bryant A, Kehoe S, Morrison J. Neoadjuvant chemotherapy before surgery versus surgery followed by chemotherapy for initial treatment in advanced ovarian epithelial cancer. *Cochrane Database Syst Rev*. 2021;7(7):CD005343.
104. van Driel WJ, Koole SN, Sonke GS. Hyperthermic intraperitoneal chemotherapy in ovarian cancer. *N Engl J Med*. 2018;378(14):1363-1364.
105. Aronson SL, Lopez-Yurda M, Koole SN, et al. Cytoreductive surgery with or without hyperthermic intraperitoneal chemotherapy in patients with advanced ovarian cancer (OVHIPEC-1): final survival analysis of a randomised, controlled, phase 3 trial. *Lancet Oncol*. 2023;24(10):1109-1118. doi:10.1016/S1470-2045(23)00396-0
106. Lim MC, Chang SJ, Park B, et al; HIPEC for Ovarian Cancer Collaborators. Survival after hyperthermic intraperitoneal chemotherapy and primary or interval cytoreductive surgery in ovarian cancer: a randomized clinical trial. *JAMA Surg*. 2022;157(5):374-383. doi:10.1001/jamasurg.2022.0143
107. Antonio CC, Alida GG, Elena GG, et al. Cytoreductive surgery with or without HIPEC after neoadjuvant chemotherapy in ovarian cancer: a phase 3 clinical trial. *Ann Surg Oncol*. 2022;29(4):2617-2625. doi:10.1245/s10434-021-11087-7

108. McGuire WP, Hoskins WJ, Brady MF, et al. Cyclophosphamide and cisplatin compared with paclitaxel and cisplatin in patients with stage III and stage IV ovarian cancer. *N Engl J Med*. 1996;334(1):1-6. doi:10.1056/NEJM199601043340101
109. Piccart MJ, Bertelsen K, James K, et al. Randomized intergroup trial of cisplatin-paclitaxel versus cisplatin-cyclophosphamide in women with advanced epithelial ovarian cancer: three-year results. *J Natl Cancer Inst*. 2000;92(9):699-708. doi:10.1093/jnci/92.9.699
110. Liu S, Kasherman L, Fazelzad R, et al. The use of bevacizumab in the modern era of targeted therapy for ovarian cancer: a systematic review and meta-analysis. *Gynecol Oncol*. 2021;161(2):601-612. doi:10.1016/j.ygyno.2021.01.028
111. Li N, Zhu J, Yin R, et al. Treatment with niraparib maintenance therapy in patients with newly diagnosed advanced ovarian cancer: a phase 3 randomized clinical trial. *JAMA Oncol*. 2023;9(9):1230-1237. doi:10.1001/jamaoncol.2023.2283
112. Monk BJ, Parkinson C, Lim MC, et al. A randomized, phase III trial to evaluate rucaparib monotherapy as maintenance treatment in patients with newly diagnosed ovarian cancer (ATHENA-MONO/GOG-3020/ENGOT-OV45). *J Clin Oncol*. 2022;40(34):3952-3964. doi:10.1200/JCO.22.01003
113. Coleman RL, Fleming GF, Brady MF, et al. Veliparib with first-line chemotherapy and as maintenance therapy in ovarian cancer. *N Engl J Med*. 2019;381(25):2403-2415. doi:10.1056/NEJMoa1909707
114. Torre LA, Trabert B, DeSantis CE, et al. Ovarian cancer statistics, 2018. *CA Cancer J Clin*. 2018;68(4):284-296. doi:10.3322/caac.21456
115. Baert T, Ferrero A, Sehoul J, et al. The systemic treatment of recurrent ovarian cancer revisited. *Ann Oncol*. 2021;32(6):710-725. doi:10.1016/j.annonc.2021.02.015
116. Wu XH, Zhu JQ, Yin RT, et al. Niraparib maintenance therapy in patients with platinum-sensitive recurrent ovarian cancer using an individualized starting dose (NORA): a randomized, double-blind, placebo-controlled phase III trial. *Ann Oncol*. 2021;32(4):512-521. doi:10.1016/j.annonc.2020.12.018
117. Wu X, Zhu J, Yin R, et al. Niraparib maintenance therapy using an individualised starting dose in patients with platinum-sensitive recurrent ovarian cancer (NORA): final overall survival analysis of a phase 3 randomised, placebo-controlled trial. *EClinicalMedicine*. 2024;72:102629. doi:10.1016/j.eclinm.2024.102629
118. Pignata S, Lorusso D, Joly F, et al; MITO16b/MANGO-OV2/ENGOT-OV17 Investigators. Carboplatin-based doublet plus bevacizumab beyond progression versus carboplatin-based doublet alone in patients with platinum-sensitive ovarian cancer: a randomised, phase 3 trial. *Lancet Oncol*. 2021;22(2):267-276. doi:10.1016/S1470-2045(20)30637-9
119. Moore KN, Oza AM, Colombo N, et al. Phase III, randomized trial of mirvetuximab soravtansine versus chemotherapy in patients with platinum-resistant ovarian cancer: primary analysis of FORWARD I. *Ann Oncol*. 2021;32(6):757-765. doi:10.1016/j.annonc.2021.02.017
120. Meric-Bernstam F, Makker V, Oaknin A, et al. Efficacy and safety of trastuzumab deruxtecan in patients with *HER2*-expressing solid tumors: primary results from the DESTINY-PanTumor02 phase II trial. *J Clin Oncol*. 2024;42(1):47-58. doi:10.1200/JCO.23.02005
121. National Comprehensive Cancer Network. Palliative care (version 2.2025). Accessed April 5, 2025. https://www.nccn.org/professionals/physician_gls/pdf/palliative.pdf
122. Bakitas MA, Tosteson TD, Li Z, et al. Early versus delayed initiation of concurrent palliative oncology care: patient outcomes in the ENABLE III randomized controlled trial. *J Clin Oncol*. 2015;33(13):1438-1445. doi:10.1200/JCO.2014.58.6362
123. Hoerger M, Wayser GR, Schwing G, Suzuki A, Perry LM. Impact of interdisciplinary outpatient specialty palliative care on survival and quality of life in adults with advanced cancer: a meta-analysis of randomized controlled trials. *Ann Behav Med*. 2019;53(7):674-685. doi:10.1093/abm/kay077