

Food Insecurity Interventions to Improve Blood Pressure

The Healthy Food First Factorial Randomized Clinical Trial

Seth A. Berkowitz, MD, MPH; Alice S. Ammerman, DrPH, MPH; Patricia Knoepf, MPH; Robert E. Anderson III, MS; Lori H. Taylor, BSc; Jenefer M. Jelede, PhD, MSc; Jessica Archibald, MSA; Keyi Xue, BSPH; Eleanor Wertman, MPH; Benjamin Dellva, MPH; Kevin Pignone, BSPH; Bahjat Qaqish, MBBS, PhD; Rowena J. Dolor, MD, MHS; Stephanie Turner, RN, MSN; John R. Lumpkin, MD, MPH; Darren A. DeWalt, MD, MPH

IMPORTANCE Food insecurity is associated with worse blood pressure control, but the optimal design for a food insecurity intervention to improve blood pressure is unknown.

OBJECTIVE To inform food insecurity intervention design by comparing different intervention elements: type of food resources provided, whether to offer lifestyle counseling, and intervention duration.

DESIGN, SETTING, AND PARTICIPANTS A $2 \times 2 \times 2$ factorial comparative effectiveness randomized clinical trial was carried out including adults with hypertension and systolic blood pressure (SBP) of 130 mm Hg or higher, who spoke English or Spanish, and reported food insecurity in 2 clinical networks across 364 clinical sites in North Carolina.

INTERVENTIONS Food resources included healthy food subsidy redeemable at grocery stores vs biweekly healthy food box home delivery. The lifestyle intervention included either no intervention or offering telephone-based lifestyle counseling. The intervention duration included 6 months vs 12 months.

MAIN OUTCOMES AND MEASURES The primary outcome was SBP. Secondary outcomes were diastolic blood pressure (DBP) and food security. The primary time point was 6 months from randomization. Twelve and 18 months were secondary time points.

RESULTS Overall, 458 individuals were randomized. The mean (SD) age was 49.7 (10.7) years and 345 (75.3%) were female individuals. Fewer than 11 participants identified as American Indian/Alaska Native; 11 (2.4%) identified as Asian, 237 (51.7%) identified as Black, 20 (4.4%) identified with multiple races, fewer than 11 participants identified as Native Hawaiian/Pacific Islander, 165 (36.0%) identified as White, and 22 (4.8%) did not report a racial identity. Twenty-two participants (4.8%) identified as Hispanic ethnicity. Mean (SD) preintervention SBP and DBP were 138.2 (11.9) and 87.4 (9.1) mm Hg, respectively. The food subsidy, compared with the food box, led to moderately lower SBP at the 6-month primary time point (132.8 vs 135.3 mm Hg; difference -2.5 mm Hg; 95% CI -4.1 to -0.9 ; $P = .003$). DBP was also lower at 6 months (80.5 vs 82.1 mm Hg; difference -1.5 mm Hg; 95% CI, -2.5 to -0.6). The food subsidy group also had lower SBP and DBP at 18 months (SBP difference, -2.1 mm Hg; 95% CI, -4.2 to -0.05 ; DBP difference, -1.6 mm Hg; 95% CI -2.8 to -0.3). SBP and DBP differences at 12 months were in favor of the food subsidy, but not significantly different. Offering lifestyle counseling did not produce significantly lower SBP or DBP than not offering counseling at any time point. The 12-month duration did not produce significantly lower SBP or DBP than 6-month duration at any time point. 6-, 12-, and 18-month food security scores decreased from baseline in all groups, and did not differ significantly between groups.

CONCLUSIONS AND RELEVANCE In this randomized comparative effectiveness trial, a food subsidy produced a moderate reduction in SBP and DBP compared with a delivered food box. Offering lifestyle counseling and a longer benefit duration did not produce better blood pressure outcomes. Food insecurity declined from baseline in all groups, but did not differ between groups.

TRIAL REGISTRATION ClinicalTrials.gov Identifier: [NCT05048836](https://clinicaltrials.gov/ct2/show/study/NCT05048836)

JAMA Intern Med. doi:10.1001/jamainternmed.2025.5287
Published online October 13, 2025.

[+ Visual Abstract](#)

[+ Editor's Note](#)

[+ Supplemental content](#)

Author Affiliations: Author affiliations are listed at the end of this article.

Corresponding Author: Seth A. Berkowitz, MD, MPH, Division of General Medicine and Clinical Epidemiology, Department of Medicine, University of North Carolina at Chapel Hill School of Medicine, 5034 Old Clinic Bldg, CB# 7110, Chapel Hill, NC 27599 (seth_berkowitz@med.unc.edu).

Food insecurity, uncertain access to the food needed for an active, healthy life,¹ affected more than 47 million people in the US in 2023.¹ Food insecurity is associated with numerous poor health outcomes, including greater prevalence and worse control of cardiometabolic conditions like hypertension, type 2 diabetes, congestive heart failure, and chronic kidney disease.²⁻¹¹

High blood pressure is a key risk factor for cardiovascular morbidity and mortality, affecting half of all adults, yet more than 75% of people in the US with hypertension have higher blood pressure than recommended despite treatment.^{12,13} Food insecurity can worsen blood pressure through at least 3 pathways.^{14,15} First, through a nutrition pathway, it can incentivize consumption of more affordable but less healthful foods. Second, through a compensatory pathway, it can force individuals to make trade-offs between food, medications, and other items necessary for good health. Finally, through a psychological pathway, it can increase stress, anxiety, and depressive symptoms and worsen overall quality of life, all of which are known to increase blood pressure.

Given the health impacts of food insecurity, there is little doubt that addressing it improves health in general. However, there are many unanswered questions regarding the effect of specific versions of food insecurity interventions on blood pressure outcomes.

To help inform how to design a food insecurity intervention that could be implemented as an insurance benefit, we conducted a pragmatic, factorial comparative effectiveness randomized clinical trial. We examined 3 dimensions of treatment in which there are unanswered questions. The first dimension was the modality of food resources provided—either a food subsidy individuals could use to purchase healthy foods of their choice at local supermarkets, or a home-delivered food box containing fresh produce and other healthy food items such as olive oil and nuts. The second dimension was lifestyle—either an offer to receive telephone-based implementation of a previously tested lifestyle intervention¹⁶⁻¹⁹ delivered by community health workers, or no offer for lifestyle counseling. The third dimension was duration—receiving the assigned intervention for 6 months or 12 months. For each dimension, we expected that the more resource-intensive intervention (food box, offer of lifestyle counseling, and 12 months duration, respectively) would lead to lower systolic and diastolic blood pressure and lower food insecurity than the less resource-intensive intervention.

Methods

Study Design

The trial protocol is available in [Supplement 1](#). This was a $2 \times 2 \times 2$ full factorial pragmatic randomized clinical trial designed to compare the effectiveness of different dimensions of food insecurity interventions meant to improve blood pressure.²⁰⁻²² The objective was to generate information that informs how to structure an insurance benefit to address food insecurity. The rationale for a factorial trial was to efficiently evaluate several dimensions of food insecurity interventions

Key Points

Question When designing a food insecurity intervention to improve blood pressure, does a food subsidy lead to better control than a food box; does the offer of lifestyle counseling improve outcomes relative to not offering lifestyle counseling; and does longer duration of intervention improve outcomes relative to shorter duration?

Findings In this randomized comparative effectiveness trial including 458 participants, a food subsidy produced better blood pressure control than a delivered food box. Offering lifestyle counseling and a longer benefit duration did not produce better blood pressure outcomes.

Meaning A food subsidy improved blood pressure control compared with a food box, but offering lifestyle counseling or longer intervention duration did not lead to better outcomes.

simultaneously. We did not design the study with the anticipation of synergistic interactions between components, but our analytic strategy did allow for the possibility of interactions. The study followed the Consolidated Standards of Reporting Trials ([CONSORT](#)) reporting guidelines.

Recruitment, Eligibility, Enrollment, and Randomization

Participants in this study were enrolled from 2 health care clinical networks in central North Carolina. Eligible participants were adults (age ≥ 18 years) who had a diagnosis of hypertension and a systolic blood pressure of 130 mm Hg or higher in the past year, spoke English or Spanish, and reported food insecurity on the 2-item Hunger Vital Sign.²³ Recruitment began in October 2021, the last participant was enrolled in January 2023, and follow-up data collection ended in August 2024. Participants provided verbal informed consent. The University of North Carolina at Chapel Hill institutional review board approved this study. This trial was registered on ClinicalTrials.gov as [NCT05048836](#) prior to enrollment of the first participant. After providing informed consent and completing baseline assessment, participants were randomly assigned to 1 of 8 treatment arms in a 1:1:1:1:1:1:1:1 ratio. A study team member not involved with enrollment prepared randomization tables using a computer-generated random number sequence prior to the first participant's enrollment, which were concealed from all other study staff using the randomization function within REDCap.²⁴ Randomization used a permuted block algorithm with blocks of size 8 and 16 in unpredictable order, and was stratified by sex (female or not), race and ethnicity (non-Hispanic White or not), and preintervention systolic blood pressure (>150 mm Hg or not). Analyses were conducted from October 2024 to April 2025.

Intervention

This factorial trial had 3 dimensions of interventions, with an individual assigned to 1 of 2 treatment conditions for each dimension, resulting in 8 possible combinations of treatment conditions. These dimensions were food resources, lifestyle, and duration. For each dimension, 1 intervention was intentionally less resource-intensive than the other and was considered the reference arm.

Along the food resources dimension, a participant could be assigned to either a food subsidy intervention or a food box intervention. The food subsidy intervention consisted of a \$40-monthly food subsidy loaded onto an electronic card that could be used to purchase fruits, vegetables, nuts, or legumes with no added sugar, salt, or fat at a local grocery store chain. Unused funds carried over from month to month until the end of the intervention. The food box arm consisted of home delivery, every other week, of a healthy food box containing produce and alternately a prepared meal or 1 dozen eggs. At the start of the intervention, participants received a stock up box with no-salt seasonings and healthy fats such as olive oil and nuts. The approximate value in the food box arm was \$115-monthly (\$100 for food, \$15 for delivery costs). The food subsidy was the reference arm for this dimension.

Along the lifestyle dimension, participants could be assigned to either be offered or not be offered telephone-based lifestyle counseling focused on improving diet quality delivered by health care system-based community health workers. The curriculum was an adaptation of the Med-South Lifestyle Program, which focuses on achieving heart-healthy Mediterranean diet patterns using foods that are affordable and familiar to individuals living in the southeastern US.^{16-19,25-28} Participants offered the counseling received a printed version of this curriculum, which included both healthy recipes and links to videos demonstrating meal preparation. The Med-South Lifestyle Program has demonstrated both high acceptability and efficacy in achieving diet quality improvements in prior randomized studies.^{16-19,25-27} The intervention consisted of monthly sessions (eTable 1 in [Supplement 2](#)). Not receiving an offer for lifestyle counseling was the reference arm for this dimension.

Along the duration dimension, individuals could be assigned to receive their intervention (defined along the other 2 dimensions) for either 6 or 12 months. The 6-month arm was the reference arm for this dimension.

Outcomes and Data Collection

The primary outcome was systolic blood pressure (SBP), recorded in routine outpatient care and extracted from the clinical data warehouses of the participating health care systems. Diastolic blood pressure (DBP) was extracted similarly. As pragmatic assessments, blood pressure observations were made during routine care, rather than on a set schedule. Those measuring the blood pressure in clinics were generally unaware of an individual's participation in the study or study arm.

A second stream of data collection came from surveys completed by participants at 4 time points: baseline (prior to randomization), and 6, 12, and 18 months after randomization. These surveys collected participant-reported information not available in the clinical data warehouse. The main outcome for these surveys was food insecurity, measured using the 10-item US Department of Agriculture (USDA) Household Food Security Survey Module with a 30-day look-back period.²⁹ Other outcomes included health-related quality of life as indicated by the Patient-Reported Outcomes Measurement Information System (PROMIS) preference (PROPr) score³⁰⁻³² (which indicates overall health-related quality of life and has subdo-

main of cognitive function, depressive symptoms, fatigue, pain interference, physical function, ability to participate in social roles, and sleep), and indicators of cost-related medication underuse and trade-offs between food and medications.^{33,34} To help understand mechanisms whereby intervention dimensions may have produced effects (particularly for the lifestyle dimension), we also measured diet quality, diet self-efficacy, and diet self-perception.^{16,35-39} Telephone surveys were administered by the Carolina Survey Research Lab, and those administering the surveys were unaware of study arm assignment while collecting outcome measures.

Statistical Analysis

For sample size determination, we targeted having 80% or greater power to detect a difference in systolic blood pressure of 3 mm Hg or greater for analyses along 1 dimension of intervention (eg, comparing food subsidy to food box groups). A sample size of 280 total participants was sufficient to provide this level of power, and enrolling a sample of this size was our primary recruitment goal. To have the same power to test hypotheses by groups defined along all 3 dimensions of intervention (eg, comparing the food subsidy with not offering lifestyle counseling for 6 months duration to the food box with offering lifestyle counseling for 12 months groups), a sample size of 1400 would have been required. We were unable to recruit a sample of 1400 owing to slower than anticipated recruitment during the COVID-19 public health emergency. The study team made the decision to halt enrollment prior to reaching this higher number of participants based on rates of enrollment and the time available to complete study activities.

We used an intention-to-treat analytic approach, analyzing all randomized participants as part of the treatment arms to which they were assigned. The estimand of interest was an average treatment effect comparing one treatment arm to another (eg, food subsidy compared with food box) while standardizing over the other treatment arms. We prespecified that all models would include interaction terms between treatment dimensions, and so all estimates allow for the possibility of interactions between treatment arms.

For blood pressure analyses (statistical analysis in [Supplement 2](#); eTable 24, model output in [Supplement 3](#)), we fit models with indicators for each treatment arm and interactions between the 3 dimensions. To help increase precision, we included the randomization stratification variables, a baseline measure of the outcome (eg, SBP for SBP analyses), and indicators of health insurance and health care system. We modeled time relative to baseline flexibly using splines with knots at 6 and 12 months after baseline, which allowed the relationship between treatment arm and time to vary at key study time points.⁴⁰⁻⁴² We fit linear models with generalized least squares, with a first-order continuous autoregressive correlation structure within subject because measurements closer together in time may be more similar than measurements further apart in time.⁴³ This longitudinal modeling approach is robust to missing data within individuals under the equivalent of a missing at random assumption,⁴⁴ so we did not implement an additional missing data strategy for these analyses. After fitting the models, we used least-squares means (also called "marginal

effects”) to create estimates at key study time points (6 [primary], 12 [secondary], and 18 [secondary] months), and contrasted them.^{45,46} The purpose of the 18-month time point was to examine the durability of intervention effects.

For analyses of the participant-reported outcomes, we used a similar approach, except we fit models using generalized estimating equations, which can accommodate binary outcomes.^{43,47} To account for missing data in these analyses, we used multiple imputation with chained equations, creating 100 imputed datasets, conducting analyses in each of them, and combining estimates using Rubin rules.⁴⁸⁻⁵⁰

We considered the SBP outcome for the food dimension at the 6-month time point to be the primary outcome and time point for type I error purposes, and report a *P* value for this analysis. We used a statistical significance threshold of $<.05$ with a 2-sided test. For other analyses, we report 95% CIs. Because we did not conduct global comparisons (eg, at least 1 of the more intensive treatment conditions is superior to the least intensive treatment condition), we did not implement adjustments for family-wise error.⁵¹⁻⁵³ Analyses were conducted in SAS statistical software (version 9.4; SAS Institute, Inc) and R statistical software (version 4.3.1; R Foundation). Statistical code for all models was developed with the analysts masked to treatment assignments, and data were only unmasked after the analytic code was finalized.

Results

Overall, 2792 individuals were assessed, 525 enrolled, and 458 were randomized after completing baseline assessments; 67 dropped out prior to randomization (**Figure 1**; eTable 2 in [Supplement 1](#)). The 458 randomized participants formed the intention-to-treat cohort that was analyzed.

The mean (SD) age of those randomized was 49.7 (10.7) years and 345 (75.3%) were female individuals. Fewer than 11 participants identified as American Indian/Alaska Native; 11 (2.4%) identified as Asian, 237 (51.7%) identified as Black, 20 (4.4%) identified with multiple races, fewer than 11 participants identified as Native Hawaiian/Pacific Islander, 165 (36.0%) identified as White, and 22 (4.8%) did not report a racial identity. Twenty two participants (4.8%) identified as Hispanic ethnicity. The mean (SD) SBP in the year prior to randomization was 138.2 (11.9) mm Hg and mean (SD) DBP was 84.8 (9.1) mm Hg (**Table 1**; eTables 3-7 in [Supplement 2](#)). Participants came from 2 clinical networks (421 from the UNC health system, 37 from the Duke health system) and data were observed across a total of 364 clinical sites within those 2 networks. In this low-risk study, participants did not report any adverse events.

For the food dimension, participants redeemed a median 91% (25th percentile: 65%, 75th percentile: 100%) of the available food subsidy funds, and participants received a median 100% (25th percentile: 96%, 75th percentile: 100%) of the available food boxes. For the lifestyle dimension, participants attended a median 17% (25th percentile: 8%, 75th percentile: 50%) of available sessions (eTable 8 in [Supplement 2](#)).

For the food dimension, the mean SBP and DBP for the food subsidy arm at 6 months was SBP: 132.8 (95% CI, 131.6-134.0) mm Hg and DBP: 80.5 (95% CI, 79.8-81.2) mm Hg. For the food box at 6 months, mean SBP was 135.3 (95% CI, 134.1-136.4) mm Hg and mean DBP was 82.1 (95% CI, 81.4-82.8) mm Hg (**Figure 2**; **Table 2**; eTable 9 in [Supplement 2](#)). The food box arm had significantly higher SBP (difference in means, 2.5 mm Hg; 95% CI, 0.9-4.1; *P* = .003) than the food subsidy arm. DBP was also higher (difference in means, 1.6 mm Hg; 95% CI, 0.6-2.5). Results were similar at 18 months, but the difference between arms was smaller at 12 months. There were no meaningful differences between the food subsidy and food box with regard to food security (**Figure 3**), diet self-efficacy, diet perception, diet quality, and health-related quality of life (eTables 10-11 in [Supplement 2](#)).

For the lifestyle dimension, study outcomes were similar between the groups that were and were not offered lifestyle counseling at all time points. For the duration dimension, study outcomes were similar between the groups that received 12 and 6 months of intervention at all time points. Results by 2-dimension and 3-dimension combinations of study arms are presented in eTables 12 to 23 in [Supplement 2](#).

Discussion

In this pragmatic randomized comparative effectiveness trial, we found that a food subsidy led to lower blood pressure than a home-delivered food box. We did not find that offering lifestyle counseling improved outcomes relative to not offering it, though actual use of counseling was low, which makes results for this dimension somewhat inconclusive. We did not find that providing 12 months of food insecurity intervention led to lower blood pressure than providing 6 months. Of note, all study arms saw an average decrease in blood pressure and food insecurity, which could represent a common effect of the food insecurity interventions or regression to the mean.

Why food subsidies led to lower blood pressures than food boxes despite similar changes in food insecurity is an interesting question. Prior theoretical work suggests 2 possible explanations. First, food security is likely necessary but insufficient for improving diet.⁵⁴ That is, although it is difficult to improve diet when food insecure, increasing food security does not inherently improve diet—the key pathway to lower blood pressure in this case.⁵⁵ Second, although all food insecurity interventions address food insecurity—that is not all they do. Different food insecurity interventions may have different impacts in other areas relevant for blood pressure. For instance, the greater flexibility that food subsidies provided may have allowed participants to consume healthy diets more in accord with their preferences, even though both interventions made healthy food more available.

This study extends prior literature on food insecurity and high blood pressure. Prior studies have found that food insecurity is associated with worse blood pressure control and adverse cardiovascular outcomes for which increased blood pressure is a key mechanism.²⁻⁷ However, how best to address food insecurity to improve blood pressure was not clear. This study adds to that literature by suggesting that a food subsidy can

Figure 1. CONSORT Diagram




Table 1. Demographic Characteristics of Study Sample

Characteristic	Overall (n = 458)	Dimension, No. (%)				Length	
		Food subsidy (n = 230)	Food box (n = 228)	Lifestyle Not offered counseling (n = 231)	Offered counseling (n = 227)	6 mo (n = 230)	12 mo (n = 228)
Age at randomization, mean (SD), y	49.7 (10.7)	48.2 (11.8)	51.2 (9.2)	49.6 (10.5)	49.8 (11.0)	50.3 (10.7)	49.0 (10.7)
Sex							
Female	345 (75.3)	172 (74.8)	173 (75.9)	173 (74.9)	172 (75.8)	173 (75.2)	172 (75.4)
Male	112 (24.5)	57 (24.8)	55 (24.1)	57 (24.7)	55 (24.2)	56 (24.3)	56 (24.6)
Not reported	NA ^a	NA ^a	NA ^a	NA ^a	NA ^a	NA ^a	NA ^a
Race							
American Indian/ Alaska Native	NA ^a	NA ^a	NA ^a	NA ^a	NA ^a	NA ^a	NA ^a
Asian	11 (2.4)	NA ^a	NA ^a	NA ^a	NA ^a	NA ^a	NA ^a
Black	237 (51.7)	118 (51.3)	119 (52.2)	117 (50.6)	120 (52.9)	123 (53.5)	114 (50.0)
Multiple races	20 (4.4)	12 (5.2)	8 (3.5)	13 (5.6)	7 (3.1)	10 (4.3)	10 (4.4)
Native Hawaiian/ Pacific Islander	NA ^a	NA ^a	NA ^a	NA ^a	NA ^a	NA ^a	NA ^a
Not reported	22 (4.8)	12 (5.2)	10 (4.4)	9 (3.9)	13 (5.7)	8 (3.5)	14 (6.1)
White	165 (36.0)	81 (35.2)	84 (36.8)	83 (35.9)	82 (36.1)	83 (36.1)	82 (36.0)
Ethnicity							
Hispanic	22 (4.8)	12 (5.2)	10 (4.4)	10 (4.3)	12 (5.3)	11 (4.8)	11 (4.8)
Not Hispanic or Latino	435 (95.0)	217 (94.3)	218 (95.6)	221 (95.7)	214 (94.3)	218 (94.8)	217 (95.2)
Not reported	NA ^a	NA ^a	NA ^a	NA ^a	NA ^a	NA ^a	NA ^a
Annual household income, mean (SD), \$	42 473 (30 477)	42 990 (31 708)	41 965 (29 280)	42 223 (29 034)	42 729 (31 956)	40 819 (28 808)	44 141 (32 053)
UNC health system	421 (91.9)	212 (92.2)	209 (91.7)	214 (92.6)	207 (91.2)	215 (93.5)	206 (90.4)
Commercial health insurance	414 (90.4)	208 (90.4)	206 (90.4)	213 (92.2)	201 (88.5)	205 (89.1)	209 (91.7)
Received SNAP in last year	139 (30.5)	73 (32.0)	66 (28.9)	67 (29.3)	72 (31.7)	65 (28.3)	74 (32.7)
Receiving SNAP at time of enrollment	101 (22.1)	57 (25.0)	44 (19.3)	43 (18.8)	58 (25.6)	49 (21.3)	52 (23.0)
Monthly SNAP benefit, mean (SD), \$	255.83 (168.73)	261.41 (180.36)	249.83 (156.42)	259.30 (176.71)	252.61 (162.17)	257.72 (180.34)	254.18 (159.11)
Preintervention systolic blood pressure, mean (SD), mm Hg	138.19 (11.88)	138.42 (12.41)	137.96 (11.35)	137.90 (11.65)	138.49 (12.14)	138.13 (11.39)	138.25 (12.38)
Preintervention diastolic blood pressure, mean (SD), mm Hg	84.75 (9.11)	84.87 (9.09)	84.62 (9.15)	85.05 (9.53)	84.43 (8.67)	84.85 (8.82)	84.64 (9.42)
Food insecure ^b	361 (79.2)	181 (79.4)	180 (78.9)	180 (78.6)	181 (79.7)	182 (79.1)	179 (79.2)
Food security score, mean (SD)	4.91 (2.73)	4.98 (2.75)	4.83 (2.71)	4.86 (2.75)	4.95 (2.71)	4.99 (2.77)	4.82 (2.68)
Diet self-efficacy score, mean (SD) ^c	23.54 (3.01)	23.57 (3.08)	23.51 (2.94)	23.76 (2.88)	23.32 (3.13)	23.48 (3.03)	23.60 (3.00)
Diet perception score, mean (SD) ^d	6.12 (1.60)	6.16 (1.61)	6.08 (1.59)	6.07 (1.57)	6.17 (1.63)	6.07 (1.58)	6.17 (1.62)
Diet quality score, mean (SD) ^e	5.74 (1.92)	5.70 (1.99)	5.78 (1.85)	5.86 (1.93)	5.62 (1.90)	5.78 (1.93)	5.70 (1.92)
Put off buying food to afford medications	204 (44.7)	98 (43.0)	106 (46.5)	97 (42.4)	107 (47.1)	107 (46.5)	97 (42.9)
Put off medications to afford food	190 (41.7)	88 (38.6)	102 (44.7)	92 (40.2)	98 (43.2)	100 (43.5)	90 (39.8)
PROPr HRQoL score, mean (SD) ^f	0.37 (0.18)	0.37 (0.19)	0.37 (0.18)	0.39 (0.18)	0.35 (0.19)	0.36 (0.18)	0.38 (0.19)

Abbreviations: NA, not applicable; PROPr HRQoL, PROMIS-Preference Health Related Quality of Life instrument; SNAP, Supplemental Nutrition Assistance Program; UNC, University of North Carolina.

^a Actual count is fewer than 11 and has been masked for low cell sizes.

^b All study participants screened positive for food insecurity on the 2-item hunger vital sign prior to enrollment. Results for this row report food security status as assessed at study baseline with a more detailed food insecurity instrument.

^c Diet self-efficacy scores range from 7 to 28 with higher scores indicating greater self-efficacy.

^d Diet perception scores range from 3 to 12 with lower scores indicating that a greater match between a respondent's perception of a healthy diet and nutritional recommendations.


^e Diet quality scores range from 0 to 12 with higher scores indicating greater diet quality.

^f PROPr HRQoL scores range from 0.0 to 1.0 with 1.0 indicating perfect health. The overall PROPr score is created from subscores in 7 different domains: cognitive, depressive symptoms, fatigue, pain interference, sleep, ability to participate in social roles, and physical health.

lower blood pressure, compared with a food box. At the population level, differences in systolic blood pressure of the magnitude seen in this study have been associated with long-

term benefits.⁵⁶ For example, Hardy et al⁵⁶ demonstrated that a 2-mm Hg decrease in SBP would be expected to reduce the rate of coronary heart disease by 27 per 100 000 person-

Figure 2. Blood Pressure Results



The mean systolic and diastolic blood pressure, with standard error bars, by treatment at baseline, 6-month, 12-month, and 18-month time points. Results are from marginal models with an autoregressive correlation structure for repeated observations within individuals and including the following covariates as fixed effects: food dimension assignment, lifestyle dimension assignment,

duration dimension assignment, all 2-way and 3-way interaction terms for treatment assignments, time, sex indicator, race and ethnicity indicator, high baseline blood pressure indicator, clinical network indicator, insurance type indicator, and mean blood pressure (systolic or diastolic, depending on the model's outcome) in the baseline period.

Table 2. Mean Blood Pressure and Difference in Means Between Groups^a

	Time point								
Dimension	6 mo			12 mo			18 mo		
Food	Food subsidy	Food box	Difference (95% CI)	Food subsidy	Food box	Difference (95% CI)	Food subsidy	Food box	Difference (95% CI)
SBP	132.8	135.3	2.5 (0.9 to 4.1)	133.2	133.5	0.3 (−1.8 to 2.5)	131.7	133.9	2.2 (0.1 to 4.3)
DBP	80.5	82.1	1.6 (0.6 to 2.5)	81.3	82.1	0.8 (−0.5 to 2.1)	80.7	82.1	1.5 (0.3 to 2.7)
Lifestyle	Not offered counseling	Offered counseling	Difference (95% CI)	Not offered counseling	Offered counseling	Difference (95% CI)	Not offered counseling	Offered counseling	Difference (95% CI)
SBP	133.5	134.7	1.2 (−0.5 to 2.8)	133.5	133.3	−0.2 (−2.3 to 2.0)	133.4	132.4	−1.0 (−3.1 to 1.0)
DBP	80.9	81.8	0.9 (0.0 to 1.9)	81.9	81.7	−0.1 (−1.4 to 1.1)	81.4	81.5	0.1 (−1.2 to 1.3)
Duration	6 mo	12 mo	Difference (95% CI)	6 mo	12 mo	Difference (95% CI)	6 mo	12 mo	Difference (95% CI)
SBP	134.8	133.2	−1.6 (−3.2 to 0.1)	133.1	133.7	0.6 (−1.5 to 2.8)	133.7	132.0	−1.7 (−3.8 to 0.3)
DBP	81.3	81.4	0.1 (−0.8 to 1.1)	80.9	82.8	1.9 (0.7 to 3.2)	81.6	81.3	−0.3 (−1.5 to 1.0)

Abbreviations: DBP, diastolic blood pressure; SBP, systolic blood pressure.

^a Blood pressure values, reported in mm Hg, represent least squares mean estimates from marginal models with an autoregressive correlation structure for repeated observations within individuals and including the following covariates as fixed effects: food dimension assignment, lifestyle dimension


assignment, duration dimension assignment, all 2-way and 3-way interaction terms for treatment assignments, time, sex indicator, race and ethnicity indicator, high baseline blood pressure indicator, clinical network indicator, insurance type indicator, and mean blood pressure (systolic or diastolic, depending on the model's outcome) in the baseline period.

years among non-Hispanic Black individuals and by 18 per 100 000 person-years among non-Hispanic White individuals. Had there been an arm with no food insecurity intervention, there may have been a greater SBP differential, as prior studies have found,⁵⁷ but leaving food insecurity unaddressed would have exposed participants to adverse conse-

quences given the clear connection between food insecurity and other poor health outcomes.^{2,5}

One unexpected finding was that the difference in blood pressure between the food subsidy and food box arms at 12 months was numerically different from the estimates at 6 months. The reasons for this are unclear. The study did occur during a period

Figure 3. Food Insecurity Results



The mean food security score, with standard error bars, by treatment at baseline, 6-month, 12-month, and 18-month time points. Results are from generalized estimating equation models with an autoregressive correlation structure for repeated observations within individuals and including the following covariates as fixed effects: food dimension assignment, lifestyle

dimension assignment, duration dimension assignment, all 2-way and 3-way interaction terms for treatment assignments, time, sex indicator, race and ethnicity indicator, high baseline blood pressure indicator, clinical network indicator, insurance type indicator, and baseline food security score.

of high food price inflation, and so the real value of the subsidy may have eroded over time. Alternatively because the 95% CI for the 12-month period did contain the point estimates for the other time points, it may simply have been chance variation.

This study has several implications, although we caution that these are presently speculative and future research should investigate these ideas more fully. First, low adherence is a likely explanation for the lifestyle counseling intervention's lack of benefit, especially since similar programs have been proven effective in studies that specifically recruited people interested in lifestyle intervention.^{16-19,25-27} Participants in this food insecurity-focused study may have been, on average, less interested in lifestyle intervention than those recruited specifically because of interest in lifestyle intervention. Second, the lack of difference between the 12- and 6-month arms may be related to the episodic nature of food insecurity; approximately 50% of individuals who experience food insecurity in 1 year do not experience it the next year.⁵⁸ As such, 6 months of support during a "rough patch" may be sufficient to mitigate some consequences of food insecurity while the individual works to change their circumstances. Of course, the finding that there was no difference on average does not necessarily mean that no individual would have done better with longer vs shorter intervention. Thus, in implementing a food insecurity intervention, it may make sense to reassess individual needs at 6 months to evaluate whether an individual is likely to benefit from further support rather than ending support at 6 months for all individuals.

Limitations

The findings of this randomized comparative effectiveness trial should be interpreted considering several limitations. First,

though the study was adequately powered to test hypotheses along 1 dimension of intervention, it was underpowered to detect interactions between multiple dimensions. However, because our analytic approach allowed for interactions, even if not statistically significant, we do not believe that this affects interpretation of the results that compare single dimensions. Second, this was a pragmatic trial with a sample that faced many barriers to participating in research. The lack of engagement with lifestyle counseling likely reflects competing demands that can limit engagement in lifestyle interventions. This is an important finding in its own right, and one that should be considered when implementing interventions outside of a research context. Prior studies using this intervention that involved individuals who enrolled in research with the explicit intent to improve their diet achieved a higher level of adherence and improved health outcomes.^{16-19,25-27} Though there was loss to follow-up, it was within the level of attrition we expected, and we used an appropriate missing data strategy to help mitigate its impact on analyses.

Conclusions

This pragmatic comparative effectiveness trial of food insecurity interventions to improve blood pressure found that a food subsidy led to lower blood pressure than a food box. There was little difference between being offered and not being offered lifestyle counseling or between having 6 vs 12 months of intervention. Both food insecurity and hypertension are important threats to health. This study suggests how these issues might be jointly addressed to improve population health.

ARTICLE INFORMATION

Accepted for Publication: August 19, 2025.

Published Online: October 13, 2025.

doi:10.1001/jamainternmed.2025.5287

Open Access: This is an open access article distributed under the terms of the [CC-BY-NC-ND](#)

License, which does not permit alteration or commercial use, including those for text and data mining, AI training, and similar technologies. © 2025 Berkowitz SA et al. *JAMA Internal Medicine*.

Author Affiliations: Division of General Medicine and Clinical Epidemiology, Department of Medicine, University of North Carolina at Chapel Hill School of

Medicine, Chapel Hill (Berkowitz, DeWalt); Cecil G. Sheps Center for Health Services Research, University of North Carolina at Chapel Hill, Chapel Hill (Berkowitz, Knoepp, Archibald, Pignone, DeWalt); Department of Nutrition, Gillings School of Global Public Health, University of North Carolina at Chapel Hill, Chapel Hill (Ammerman, Anderson);

Center for Health Promotion and Disease Prevention, University of North Carolina at Chapel Hill, Chapel Hill (Ammerman); Drivers of Health Strategy, Blue Cross and Blue Shield of North Carolina, Durham (Taylor, Jedeke, Lumpkin); Department of Biostatistics, Gillings School of Global Public Health, University of North Carolina at Chapel Hill (Xue, Qaqish, Turner); UNC Health Alliance, Morrisville, North Carolina (Wertman, Dellva); Division General Internal Medicine, Department of Medicine, Duke University School of Medicine, Durham, North Carolina (Dolor); Blue Cross and Blue Shield of North Carolina Foundation, Durham, North Carolina (Lumpkin).

Author Contributions: Author X had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Berkowitz, Ammerman, Knoepf, Taylor, Wertman, Turner, Lumpkin, DeWalt.

Acquisition, analysis, or interpretation of data: Berkowitz, Ammerman, Knoepf, Anderson, Jedeke, Archibald, Xue, Wertman, Dellva, Pignone, Qaqish, Dolor, DeWalt.

Drafting of the manuscript: Berkowitz, Ammerman, Knoepf, Anderson, Qaqish, DeWalt.

Critical review of the manuscript for important intellectual content: Ammerman, Anderson, Taylor, Jedeke, Archibald, Xue, Wertman, Dellva, Pignone, Qaqish, Dolor, Turner, Lumpkin, DeWalt.

Statistical analysis: Berkowitz, Anderson, Jedeke, Archibald, Xue, Qaqish.

Obtained funding: Ammerman, Lumpkin, DeWalt.

Administrative, technical, or material support: Ammerman, Knoepf, Anderson, Taylor, Archibald, Dellva, Pignone, Turner, Lumpkin, DeWalt.

Supervision: Berkowitz, Wertman, Pignone, Qaqish, Turner, Lumpkin, DeWalt.

Conflict of Interest Disclosures: Dr Berkowitz reported grants from Blue Cross Blue Shield North Carolina (BCBSNC) during the conduct of the study; grants from the National Institutes of Health (NIH), the North Carolina Department of Health and Human Services, the American Heart Association, the American Diabetes Association, and Feeding America, personal fees from Aspen Institute, Rockefeller Foundation, Gretchen Swanson Center for Nutrition, and Kaiser Permanente outside the submitted work; in addition, Dr Berkowitz had a patent for Johns Hopkins University Press with royalties paid for sales of the book *Equal Care: Health Equity, Social Democracy, and the Egalitarian State*. Dr Ammerman reported grants from BCBSNC (UNI-3001842-SOW8) during the conduct of the study; grants from NIH, the Centers for Disease Control (CDC), and the United States Department of Agriculture (USDA) outside the submitted work. Dr Knoepf reported grants from BCBSNC during the conduct of the study. Dr Anderson reported grants from BCBSNC during the conduct of the study. Dr Archibald reported grants from BCBSNC during the conduct of the study; grants from North Carolina Department of Health and Human Services (30-2021-017-DHB) outside the submitted work. Dr Dolor reported grants from BCBSNC (UNI-3001842-SOW8) during the conduct of the study. Dr Turner reported value-based care funded study from BCBSNC during the conduct of the study; contracts from BCBSNC with the University of North Carolina Health Alliance outside the submitted work. Dr DeWalt reported grants from

BCBSNC during the conduct of the study; grants from NIH, AHRQ, North Carolina Department of Health and Human Services, and the Arnold Foundation, and personal fees from BCBSNC physician advisory group outside the submitted work. No other disclosures were reported.

Funding/Support: This work was supported by BCBSNC (UNI-3001842-SOW8).

Role of the Funder/Sponsor: This study was designed as a partnership between investigators at BCBSNC and the University of North Carolina at Chapel Hill to answer questions relevant to providing food insecurity interventions as a supplemental health insurance benefit. Co-investigators, named as authors on the manuscript, from the funding organization participated in study design, and in identifying potentially eligible participants. Further, they participated as co-authors in revising the initial draft of manuscript for publication. The funding organization had no role in the collection and management of data or the decision to submit the manuscript for publication.

Meeting Presentation: Some results from this study were presented at the 2025 Society for General Internal Medicine meeting on May 16, 2025.

Data Sharing Statement: See [Supplement 4](#).

Additional Contributions: We gratefully acknowledge the contributions of the study participants and: Rawan Ajeen, BSPH, Department of Nutrition, Gillings School of Global Public Health, University of North Carolina at Chapel Hill; Joanna Aquino, BS, Clinical and Translational Sciences Institute, Duke University School of Medicine; Marcella Boynton, PhD, MS, North Carolina Translational and Clinical Sciences Institute, School of Medicine, University of North Carolina Chapel Hill; Mark Gwynne, DO, UNC Health Alliance; Tylah Harrison, Cecil G. Sheps Center for Health Services Research, University of North Carolina at Chapel Hill; Mark Holmes, PhD, Cecil G. Sheps Center for Health Services Research, University of North Carolina at Chapel Hill; Gary Hunt, BS, Cecil G. Sheps Center for Health Services Research, University of North Carolina at Chapel Hill; Neil Kamdar, MA, Cecil G. Sheps Center for Health Services Research, University of North Carolina at Chapel Hill; Curtis Kieler, BS, Clinical and Translational Sciences Institute, Duke University School of Medicine; Amy Lamb, BS, UNC Health Alliance; Samyuktha Nandhakumar, MS, North Carolina Translational and Clinical Sciences Institute, School of Medicine, University of North Carolina Chapel Hill; Eric Newsome, BS, UNC Health Alliance; Stephanie Pierson, MS, Cecil G. Sheps Center for Health Services Research, University of North Carolina at Chapel Hill; Christine Rabenda, MHA, UNC Health Alliance; Jordan Sharpe, BA, UNC Health Alliance; Tunde Sotunde, MD, MBA, Blue Cross Blue Shield of North Carolina; Shaun Ward, BS, UNC Health Alliance; Morris Weinberger, PhD, Gillings School of Global Public Health, University of North Carolina at Chapel Hill.

REFERENCES

1. Rabbitt MP, Reed-Jones M, Hales LJ, Burke MP. Household Food Security in the United States in 2023. Accessed October 3, 2024. <https://www.ers.usda.gov/publications/pub-details/?pubid=109895>

2. Gundersen C, Ziliak JP. Food insecurity and health outcomes. *Health Aff (Millwood)*. 2015;34(11):1830-1839. doi:10.1377/hlthaff.2015.0645
3. Berkowitz SA, Berkowitz TSZ, Meigs JB, Wexler DJ. Trends in food insecurity for adults with cardiometabolic disease in the United States: 2005-2012. *PLoS One*. 2017;12(6):e0179172. doi:10.1371/journal.pone.0179172
4. Brandt EJ, Mozaffarian D, Leung CW, Berkowitz SA, Murthy VL. Diet and food and nutrition insecurity and cardiometabolic disease. *Circ Res*. 2023;132(12):1692-1706. doi:10.1161/CIRCRESAHA.123.322065
5. Mozaffarian D, Aspry KE, Garfield K, et al; ACC Prevention of Cardiovascular Disease Section Nutrition and Lifestyle Working Group and Disparities of Care Working Group. "Food is medicine" strategies for nutrition security and cardiometabolic health equity: JACC state-of-the-art review. *J Am Coll Cardiol*. 2024;83(8):843-864. doi:10.1016/j.jacc.2023.12.023
6. Volpp KG, Berkowitz SA, Sharma SV, et al; American Heart Association. Food is medicine: a Presidential Advisory from the American Heart Association. *Circulation*. 2023;148(18):1417-1439. doi:10.1161/CIR.0000000000001182
7. Crews DC, Kuczmarski MF, Grubbs V, et al; Centers for Disease Control and Prevention Chronic Kidney Disease Surveillance Team. Effect of food insecurity on chronic kidney disease in lower-income Americans. *Am J Nephrol*. 2014;39(1):27-35. doi:10.1159/000357595
8. Berkowitz SA, Karter AJ, Corbie-Smith G, et al. Food insecurity, food "deserts," and glycemic control in patients with diabetes: a longitudinal analysis. *Diabetes Care*. Published online March 19, 2018. doi:10.2337/dc17-1981
9. Tait CA, L'Abbé MR, Smith PM, Rosella LC. The association between food insecurity and incident type 2 diabetes in Canada: a population-based cohort study. *PLoS One*. 2018;13(5):e0195962. doi:10.1371/journal.pone.0195962
10. Vercammen KA, Moran AJ, Carnethon MR, et al. Longitudinal analysis of food insufficiency and cardiovascular disease risk factors in the CARDIA study. *Am J Prev Med*. 2022;62(1):65-76. doi:10.1016/j.amepre.2021.06.020
11. Jia J, Carnethon MR, Wong M, Lewis CE, Schreiner PJ, Kandula NR. Food insecurity and incident cardiovascular disease among Black and White US individuals, 2000-2020. *JAMA Cardiol*. 2025;10(5):456-462. doi:10.1001/jamacardio.2025.0109
12. Centers for Disease Control and Prevention. Hypertension Prevalence in the U.S. | Million Hearts. Centers for Disease Control and Prevention. May 12, 2023. Accessed March 17, 2025. <https://millionhearts.hhs.gov/data-reports/hypertension-prevalence.html>
13. Whelton PK, Carey RM, Aronow WS, et al. A guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension*. Published online January 1, 2017. doi:10.1161/HYP.000000000000065
14. Te Vazquez J, Feng SN, Orr CJ, Berkowitz SA. Food insecurity and cardiometabolic conditions: a review of recent research. *Curr Nutr Rep*. 2021;10(4):243-254. doi:10.1007/s13668-021-00364-2
15. Palakshappa D, Ip EH, Berkowitz SA, et al. Pathways by which food insecurity is associated

- with atherosclerotic cardiovascular disease risk. *J Am Heart Assoc*. 2021;10(22):e021901. doi:10.1161/JAHA.121.021901
16. Keyserling TC, Samuel-Hodge CD, Pitts SJ, et al. A community-based lifestyle and weight loss intervention promoting a Mediterranean-style diet pattern evaluated in the stroke belt of North Carolina: the Heart Healthy Lenoir Project. *BMC Public Health*. 2016;16:732. doi:10.1186/s12889-016-3370-9
 17. Embree GGR, Samuel-Hodge CD, Johnston LF, et al. Successful long-term weight loss among participants with diabetes receiving an intervention promoting an adapted Mediterranean-style dietary pattern: the Heart Healthy Lenoir Project. *BMJ Open Diabetes Res Care*. 2017;5(1):e000339. doi:10.1136/bmjdr-2016-000339
 18. Keyserling TC, Sheridan SL, Draeger LB, et al. A comparison of live counseling with a web-based lifestyle and medication intervention to reduce coronary heart disease risk: a randomized clinical trial. *JAMA Intern Med*. 2014;174(7):1144-1157. doi:10.1001/jamainternmed.2014.1984
 19. Cené CW, Halladay JR, Gizlice Z, et al. A multicomponent quality improvement intervention to improve blood pressure and reduce racial disparities in rural primary care practices. *J Clin Hypertens (Greenwich)*. 2017;19(4):351-360. doi:10.1111/jch.12944
 20. Kahan BC, Hall SS, Beller EM, et al. Reporting of factorial randomized trials: extension of the CONSORT 2010 Statement. *JAMA*. 2023;330(21):2106-2114. doi:10.1001/jama.2023.19793
 21. Ciolino JD, Scholtens DM, Bonner LB. Factorial clinical trial designs. *JAMA*. 2025;333(6):532-533. doi:10.1001/jama.2024.25374
 22. Stampfer MJ, Buring JE, Willett W, Rosner B, Eberlein K, Hennekens CH. The 2 x 2 factorial design: its application to a randomized trial of aspirin and carotene in U.S. physicians. *Stat Med*. 1985;4(2):111-116. doi:10.1002/sim.4780040202
 23. Hager ER, Quigg AM, Black MM, et al. Development and validity of a 2-item screen to identify families at risk for food insecurity. *Pediatrics*. 2010;126(1):e26-e32. doi:10.1542/peds.2009-3146
 24. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform*. 2009;42(2):377-381. doi:10.1016/j.jbi.2008.08.010
 25. Donahue KE, Tillman J, Halladay JR, et al. Lessons learned from implementing health coaching in the Heart Healthy Lenoir Hypertension Study. *Prog Community Health Partnersh*. 2016;10(4):559-567. doi:10.1353/cpr.2016.0064
 26. Halladay JR, Donahue KE, Hinderliter AL, et al; Heart Healthy Lenoir Research Team. The Heart Healthy Lenoir project—an intervention to reduce disparities in hypertension control: study protocol. *BMC Health Serv Res*. 2013;13:441. doi:10.1186/1472-6963-13-441
 27. Halladay JR, Donahue KE, Cené CW, et al. The association of health literacy and blood pressure reduction in a cohort of patients with hypertension: the Heart Healthy Lenoir Trial. *Patient Educ Couns*. 2017;100(3):542-549. doi:10.1016/j.pec.2016.10.015
 28. Saraiya VM, Berkowitz SA, Johnston LF, et al. An evaluation of primary care clinician referrals for behavioral weight loss counseling. *J Prim Care Community Health*. 2024;15:21501319241308054. doi:10.1177/21501319241308054
 29. Bickel G, Nord M, Price C, Hamilton W, Cook J. Guide to Measuring Household Food Security, Revised 2000. Published online March 2000. Accessed September 17, 2018. <https://fns-prod.azureedge.net/sites/default/files/FSGuide.pdf>
 30. Hanmer J, Cella D, Feeny D, et al. Selection of key health domains from PROMIS for a generic preference-based scoring system. *Qual Life Res*. 2017;26(12):3377-3385. doi:10.1007/s11136-017-1686-2
 31. Dewitt B, Feeny D, Fischhoff B, et al. Estimation of a preference-based summary score for the patient-reported outcomes measurement information system: the PROMIS-Preference (PROPr) Scoring System. *Med Decis Making*. 2018;38(6):683-698. doi:10.1177/0272989X18776637
 32. Hanmer J, DeWalt DA, Berkowitz SA. Association between food insecurity and health-related quality of life: a nationally representative survey. *J Gen Intern Med*. 2021;36(6):1638-1647. doi:10.1007/s11606-020-06492-9
 33. Berkowitz SA, Seligman HK, Choudhry NK. Treat or eat: food insecurity, cost-related medication underuse, and unmet needs. *Am J Med*. 2014;127(4):303-310.e3. doi:10.1016/j.amjmed.2014.01.002
 34. Ippolito MM, Lyles CR, Prendergast K, Marshall MB, Waxman E, Seligman HK. Food insecurity and diabetes self-management among food pantry clients. *Public Health Nutr*. 2017;20(1):183-189. doi:10.1017/S1368890016001786
 35. Abernethy ES, Collins KK, Hinderliter JM, et al. Validity and reliability of a brief dietary assessment questionnaire in a cardiac rehabilitation program. *J Cardiopulm Rehabil Prev*. 2020;40(4):280-283. doi:10.1097/HCR.0000000000000505
 36. Papadaki A, Johnson L, Toumpakari Z, et al. Validation of the English version of the 14-Item Mediterranean Diet Adherence Screener of the PREDIMED Study, in people at high cardiovascular risk in the UK. *Nutrients*. 2018;10(2):138. doi:10.3390/nu10020138
 37. Martínez-González MA, García-Arellano A, Toledo E, et al; PREDIMED Study Investigators. A 14-item Mediterranean diet assessment tool and obesity indexes among high-risk subjects: the PREDIMED trial. *PLoS One*. 2012;7(8):e43134. doi:10.1371/journal.pone.0043134
 38. Pinard CA, Uvena LM, Quam JB, Smith TM, Yaroch AL. Development and testing of a revised cooking matters for adults survey. *Am J Health Behav*. 2015;39(6):866-873. doi:10.5993/AJHB.39.6.14
 39. Kraschnewski JL, Gold AD, Gizlice Z, et al. Development and evaluation of a brief questionnaire to assess dietary fat quality in low-income overweight women in the southern United States. *J Nutr Educ Behav*. 2013;45(4):355-361. doi:10.1016/j.jneb.2012.10.008
 40. Egbewale BE, Lewis M, Sim J. Bias, precision and statistical power of analysis of covariance in the analysis of randomized trials with baseline imbalance: a simulation study. *BMC Med Res Methodol*. 2014;14(1):49. doi:10.1186/1471-2288-14-49
 41. Wang B, Ogburn EL, Rosenblum M. Analysis of covariance in randomized trials: More precision and valid confidence intervals, without model assumptions. *Biometrics*. 2019;75(4):1391-1400. doi:10.1111/biom.13062
 42. Holmberg MJ, Andersen LW. Adjustment for baseline characteristics in randomized clinical trials. *JAMA*. 2022;328(21):2155-2156. doi:10.1001/jama.2022.21506
 43. Pekár S, Brabec M. Marginal models via GLS: a convenient yet neglected tool for the analysis of correlated data in the behavioural sciences. *Ethology*. 2016;122(8):621-631. doi:10.1111/eth.12514
 44. Gabrio A, Plumpton C, Banerjee S, Leurent B. Linear mixed models to handle missing at random data in trial-based economic evaluations. *Health Econ*. 2022;31(6):1276-1287. doi:10.1002/hec.4510
 45. Permutt T. Do covariates change the estimand? *Stat Biopharm Res*. 2020;12(1):45-53. doi:10.1080/19466315.2019.1647874
 46. Laffin LJ, Rodman D, Luther JM, et al; Target-HTN Investigators. Aldosterone synthase inhibition with lorundrostat for uncontrolled hypertension: the Target-HTN Randomized Clinical Trial. *JAMA*. 2023;330(12):1140-1150. doi:10.1001/jama.2023.16029
 47. Zeger SL, Liang KY, Albert PS. Models for longitudinal data: a generalized estimating equation approach. *Biometrics*. 1988;44(4):1049-1060. doi:10.2307/2531734
 48. Azur MJ, Stuart EA, Frangakis C, Leaf PJ. Multiple imputation by chained equations: what is it and how does it work? *Int J Methods Psychiatr Res*. 2011;20(1):40-49. doi:10.1002/mpr.329
 49. White IR, Royston P, Wood AM. Multiple imputation using chained equations: issues and guidance for practice. *Stat Med*. 2011;30(4):377-399. doi:10.1002/sim.4067
 50. van Buuren S, Groothuis-Oudshoorn K, Vink G, et al. mice: Multivariate Imputation by Chained Equations. Published online June 5, 2023. Accessed March 8, 2024. <https://cran.r-project.org/web/packages/mice/index.html>
 51. Rothman KJ. No adjustments are needed for multiple comparisons. *Epidemiology*. 1990;1(1):43-46. doi:10.1097/00001648-199001000-00010
 52. Bender R, Lange S. Adjusting for multiple testing—when and how? *J Clin Epidemiol*. 2001;54(4):343-349. doi:10.1016/S0895-4356(00)00314-0
 53. Greenland S, Hofman A. Multiple comparisons controversies are about context and costs, not frequentism versus Bayesianism. *Eur J Epidemiol*. 2019;34(9):801-808. doi:10.1007/s10654-019-00552-z
 54. Berkowitz SA, Seligman HK, Mozaffarian D. A new approach to guide research and policy at the intersection of income, food, nutrition, and health. *Health Aff (Millwood)*. 2025;44(4):384-390. doi:10.1377/hlthaff.2024.01346
 55. Pilla SJ, Yeh HC, Mitchell CM, et al. Dietary patterns, sodium reduction, and blood pressure in type 2 diabetes: the DASH4D randomized clinical trial. *JAMA Intern Med*. Published online June 9, 2025. doi:10.1001/jamainternmed.2025.1580
 56. Hardy ST, Loehr LR, Butler KR, et al. Reducing the blood pressure-related burden of cardiovascular disease: impact of achievable improvements in blood pressure prevention and control. *J Am Heart Assoc*. 2015;4(10):e002276. doi:10.1161/JAHA.115.002276
 57. Lapay ER, Sytsma TM, Hutchinson HM, et al. Medically tailored grocery deliveries to improve food security and hypertension in underserved groups: a student-run pilot randomized controlled trial. *Healthcare (Basel)*. 2025;13(3):253. doi:10.3390/healthcare13030253
 58. Berkowitz SA, Seligman HK, Palakshappa D. Understanding food insecurity risk in the United States: a longitudinal analysis. *SSM Popul Health*. 2023;25:101569. doi:10.1016/j.ssmph.2023.101569